

BioBasis Manual

Conceptual design and sampling procedures of the biological monitoring programme within NuukBasic



6th edition, 2022



BioBasis Manual – Conceptual design and sampling procedures of the biological monitoring programme within NuukBasic.

6th edition, April 2022.

By Katrine Raundrup, Maia Olsen, Ida B. Dyrholm Jacobsen, Josephine Nymand, Torben Lauridsen, Paul Henning Krogh, Niels Martin Schmidt, Lotte Illeris, Helge Ro-Poulsen, Peter Aastrup

Published by Greenland Institute of Natural Resources
GL-3900 Nuuk
Greenland

Front cover illustration: Bottles for lake monitoring at Qassi-sø, Kobbefjord, just before being wrapped in tin foil. Photo by Katrine Raundrup

Preface


BioBasis Nuuk monitors key species and key processes across plant and animal populations and their interactions within the terrestrial and limnic ecosystems in Kobbefjord, Nuuk, thereby documenting the intra- and inter-annual variation, resilience, and long-term trends. Emphasis is on biodiversity, abundance and composition, phenology, reproduction, and predation as important components in the structure and functioning of the low arctic ecosystems.

Additional biotic (e.g., NDVI) and abiotic parameters (e.g., snow-melt patterns, soil moisture and water chemistry) related to biota are monitored, complementing other subprograms.

BioBasis aims at maintaining the integrity of the long core time series available from Kobbefjord, Nuuk, whilst developing the monitoring to embrace new methodologies and questions. Field protocols for monitoring elements that are no longer part of the BioBasis monitoring programme can be found in previous versions of the manual.

The BioBasis subprogram is part of a larger set of subprograms (ClimateBasis, GeoBasis, GlacioBasis and MarineBasis), which combined, is referred to as NuukBasic. NuukBasic together with corresponding programmes in ZackenbergBasic (since 1995) and DiskoBasic (since 2017) form the integrated monitoring and long-term research program on ecosystem dynamics and climate change effects and feedbacks in Greenland, Greenland Ecosystem Monitoring (GEM; www.g-e-m.dk).

GEM operates within three science themes: Climate and Cryosphere, Ecosystem feedbacks, and Biodiversity and Populations. The different subprogrammes including “Remote sensing and Modelling” make up the operational structure of GEM.



Contents

1 Introduction	6
2 Overview of monitoring elements	8
2.1 Plants.....	8
2.1.1 The NERO line	8
2.1.2 Reproductive phenology	9
2.1.3 Total flowering	9
2.1.4 Normalized Difference Vegetation Index (NDVI)	9
2.1.5 CO ₂ flux plots.....	9
2.1.6 UV-B exclusion	9
2.2 Arthropods	10
2.3 Microarthropods.....	10
2.4 Birds	10
2.5 Mammals	10
2.6 Lakes.....	10
3. Detailed manual.....	10
3.1 Plants.....	10
3.1.1 The NERO line	10
3.1.2 Reproductive phenology.....	12
3.1.3 Total flowering	20
3.1.4 Normalised Difference Vegetation Index (NDVI) in plots and along the NERO line	22
3.1.5 Normalised Difference Vegetation Index (NDVI) in CO ₂ flux plots	25
3.1.6 CO ₂ flux plots.....	27
3.1.7 UV-B exclusion	32
3.2 Arthropods	35
3.3 Microarthropods.....	39
3.4 Birds	43
3.4.1 Breeding phenology of passerines.....	43
3.4.2 Point sampling.....	43
3.5 Mammals	45
3.6 Lakes.....	45
3.6.1 Water samples	46
3.6.2 Submerged vegetation monitored annually	49
3.6.3 Parameters monitored every 5 th year.....	49
3.7 Random observations of birds and mammals	52

3.8 Disturbance	53
4 Storage of data.....	54
4.1 Data at GINR.....	54
4.2 Data for the GEM database.....	54
5 References	56
6 List of scientific and technical consultants.....	56
7 Appendix	57
Appendix 1 Example of annual sampling protocol.....	57
Appendix 2 Data form for plant phenology monitoring	58
Appendix 3 Data form for CO ₂ -flux monitoring.....	60
Appendix 4 Data form for plant phenology in CO ₂ -flux plots	61
Appendix 5 Data form for UV-B effects monitoring.....	62
Appendix 6 Data form for arthropod monitoring	63
Appendix 7 Data form for nest phenology.....	64
Appendix 8 Data form for passerine bird monitoring.....	65
Appendix 9 Data form for random observations	66
Appendix 10 Data form for lake monitoring (Badesø)	67
Appendix 11 Data form for lake monitoring (Qassi-sø)	68
Appendix 12 Data form for submerged vegetation monitoring in lakes	69
Appendix 13 Measuring chlorophyll <i>a</i> in the laboratory	70
Appendix 14 Labels for arthropod samples	71
Appendix 15 Labels for microarthropod samples	72
Appendix 16 List with random numbers for microarthropod sampling	73

1 Introduction

The NuukBasic programme was initiated in 2007 by the National Environmental Research Institute, Aarhus University, in cooperation with the Greenland Institute of Natural Resources. NuukBasic is funded by the Danish Energy Agency and the Danish Environmental Protection Agency. The present manual describes methods and sampling procedures of the subprogram BioBasis. The manual is updated regularly. The latest version can always be found at www.g-e-m.dk.

NuukBasic is a climate change effects monitoring programme with the study area located close to Nuuk in west Greenland. The programme studies the effects of climate variability and change on marine and terrestrial ecosystems. In terms of scientific concept, NuukBasic is equivalent to the investigations carried out in ZackenbergBasic, at Zackenberg Research Station in Northeast Greenland (www.zackenberg.dk).

The terrestrial monitoring area is located at the inner parts of Kobbefjord approximately 20 km east of Nuuk within the 32km² study area (figure 1 and 2). The local climate is low arctic with a mean annual temperature of - 0.1°C (2008-2020). The warmest month is July (average 10.6°C) while the average temperature in February is - 8.6°C (coldest month). The annual mean precipitation in Kobbefjord is ca. 830 mm and maximum snow depths vary strongly between 0.3 and 1.3 m at the climate station. The drainage basin is located in an alpine landscape with mountains rising up to 1400 meter above sea level and with glacier coverage of approximately 2km². Geologically, the area is relatively homogenous with precambrium gneisses as basement throughout the drainage basin. For a presentation of the Kobbefjord geology please read the note on “Kobbefjord geology” by Thebaud and Lebrun which can be found in the main cabin at the Kobbefjord Research Station.




Figure 1. The study area. The orange symbol marks the location of the Kobbefjord Research Station. Nuuk is located further east (outside the figure to the left).




Figure 2. Upper figure: Close up of the BioBasis study area. The climate station is marked by a black dot, the heath site by a green dot and the fen site by a red dot. The orange symbol marks the location of the Kobbefjord Research Station. Lower figure: Close up of area with most plots included. The two large dark green dots refer to the GeoBasis stations: Fen site and Heath site. Please note that some of the plots are very close together and points on the map may be overlapping.

2 Overview of monitoring elements

Table 1 gives an overview of the monitoring elements. GPS-positions given throughout this manual are in decimal degrees.

Table 1. Overview of monitoring elements.

Monitoring element	Species, treatment, sample	Number of plots, samples	Sampling frequency	Sampling object	Sampling period
NERO line	Plant communities and species	-	Every 5 th year		
Phenology	<i>Salix glauca</i>	4	Weekly	Buds, flowers, senescent	May-October
	<i>Loiseleuria procumbens</i>	4			
	<i>Silene acaulis</i>	4			
Total count of flowering shoots	<i>Salix glauca</i>	4	Once per season at peak flowering	Flowers (Sal, Loi, Sil), individual plants (Eri)	Depending on phenology
	<i>Loiseleuria procumbens</i>	4			
	<i>Silene acaulis</i>	4			
	<i>Eriophorum angustifolium</i>	4			
NDVI	Along NERO line	-	Monthly	Plot	May-October
	In phenology plots incl. <i>E. angustifolium</i> and <i>Empetrum nigrum</i>	20	Weekly		May-October
CO₂-flux	Control, C	6	Weekly	Plot	May-October
	Increased temperature, T	6			
	Shading, S	6			
	Long growing season, LG	6			
	Short growing season, SG	6			
	Vegetation pin-point analysis	30	Every 5 th year	Plot (inside frame)	
UV-B	Control, C	5	3 times per season	Plot	Weekly in weeks 29-31
	Mylar film, B	5			
	Teflon film, F	5			
Arthropod	All taxonomic groups	4, 20 samples	Weekly	Specimens	May-October
Microarthropod	All taxonomic groups	8, 32 samples	3 times per season	Specimens	June-September
Birds	Passerines	13	Weekly	Specimens	May-October
	Passerine nests	Ad hoc	As often as possible	Nests	Breeding season
Mammals	Arctic fox, caribou	Ad hoc	When observed	Specimens	All year
Lakes	Water chemistry, zoo- and phytoplankton, Chl <i>a</i> , DOC	2 lakes	5 times per season		Ice-free period
	Flora	2 lakes	Once per season		
	Fauna (Arctic char, sticklebacks)	1 lake	Every 5 th year		
	Isotopes	2 lakes	Every 5 th year		

2.1 Plants

2.1.1 The NERO line

A permanent vegetation transect (the NERO line) was established in July 2007 to monitor future changes in the distribution and composition of vascular plant species in the plant communities (Bay et al. 2008). Surveys of the transect take place with 5-year intervals and have been done in 2007, 2012 and 2017. In 2010, mosses and lichens were included in the monitoring programme.

The concept relies on the assumption that changes in the distribution of plant communities can be observed by tracking changes of boundary lines between vegetation zones. Thus, each boundary between vegetation zones has been marked by a peg. The species composition of the vegetation zones is documented by Raunkjær analyses. Immigration of new species is assumed to be documented by the surveys with five-year intervals. The same concept is used for the ZERO line in Zackenberg in high arctic Northeast Greenland (Fredskild & Mogensen 1996).

2.1.2 Reproductive phenology

It is expected that plant phenology will give an early and distinct response to climate change. This has already been convincingly shown in Zackenberg (Post et al. 2009). In Nuuk we monitor three species: *Salix glauca* SAL, *Loiseleuria procumbens* LOI (new accepted name: *Kalmia procumbens*), and *Silene acaulis* SIL. These species were chosen because they are widely distributed in the area, they cover a spectrum of different growth forms (deciduous dwarf shrub, evergreen dwarf shrub, and cushion forming herb), and they are functionally comparable to species monitored in Zackenberg.

For each species, four monitoring plots were established in 2007/2008. The specific sites of the plots were chosen to cover the ecological amplitude of the species with respect to duration of snow cover, difference in soil moisture and altitude. The size of each plot varies depending on the abundance of individual flowers/catkins of the species in question.

2.1.3 Total flowering

Total flowering is a proxy of maximum annual productivity. Total flowering in the reproductive phenology plots is monitored in *Salix glauca*, *Loiseleuria procumbens*, *Eriophorum angustifolium* ERI, and *Silene acaulis*. The number of flowers is counted at peak flowering as the total number of buds, flowers (catkins in *Salix* and flowering stems in *Eriophorum*) and senescent flowers (catkins in *Salix* and flowering stems in *Eriophorum*).

2.1.4 Normalized Difference Vegetation Index (NDVI)

NDVI is a measure of vegetation greenness and used as an index of plant production and vigorousness. The progression in vegetation greenness is monitored in the plant phenology plots including four plots with *Empetrum nigrum* EMP and in the CO₂ flux plots by measuring NDVI with a scanner. The scanner measures the spectral reflectance of the plant canopy. Up to and including 2016 NDVI was also monitored along the NERO line.

2.1.5 CO₂ flux plots

The CO₂ flux is important for understanding the balance between emission (respiration) and uptake (photosynthesis) of CO₂ in the vegetation. The monitoring documents the present state, but also provides data from manipulations simulating increased temperature and increased cloud cover. This will provide an indication of whether the low arctic vegetation type dwarf shrub heath functions as a sink or a source of CO₂ now as well as in the future, where increased temperatures and cloud cover is expected.

2.1.6 UV-B exclusion

UV-B radiation will increase as a result of the expected depletion of the ozone layer in the atmosphere (Jansen et al. 1998). UV-B radiation has damaging effects on higher plants as e.g., normal growth and development may be affected. We monitor the effect of increased UV-B radiation on plant stress indirectly by measuring chlorophyll fluorescence in three series of plots: Controls, plots with a filter excluding UV-B, and filter controls with a film without exclusion of UV-B. Measurements of chlorophyll fluorescence are carried out on individual *Betula nana* and *Vaccinium uliginosum* leaves in a mesic dwarf shrub heath dominated by *Empetrum nigrum* and with *Betula nana* and *Vaccinium uliginosum* as subdominant species.

2.2 Arthropods

Arthropods constitute one of the most important groups in the arctic food chain. The group contains both herbivores and predators and monitoring of them pertains to both biodiversity and community composition as well as abundance and emergence phenology. Arthropods are sampled by means of yellow pitfall traps. Window traps are used for flying insects. The traps are emptied weekly throughout the snow free season. Samples are stored at the Greenland Institute of Natural Resources (GINR) and shipped to, sorted, and identified at the Department of Ecoscience, Aarhus University, Silkeborg. Samples are stored at Naturhistorisk Museum, Aarhus.

2.3 Microarthropods

Microarthropods are major contributors to the decomposition of organic material in the soil. They are collected from soil cores three times during the snow free season using a soil auger (soil corer). The organisms are extracted in a heat extractor by gradually heating. Microarthropods are identified at Department of Ecoscience, Aarhus University, Silkeborg.

2.4 Birds

The avifauna is monitored with special emphasis on passerine birds representing the highest trophic level monitored in Kobbefjord. Breeding phenology (first egg dates, hatching, fledging) is monitored throughout the season on an ad hoc basis.

Weekly counts of birds are carried out at 13 census points during the snow free season. Other bird observations are recorded ad-hoc during the entire field season.

2.5 Mammals

Only few terrestrial mammals occur in the study area and only sporadically: Arctic fox (*Vulpes lagopus*), arctic hare (*Lepus arcticus*), and caribou (*Rangifer tarandus*). All observations of mammals are recorded ad-hoc. If arctic fox dens are discovered, reproduction will be followed.

2.6 Lakes

Three lakes are located in the Kobbefjord catchment area: Badesø / Kangerluarsunnguup Tasia (64.13°N, -51.36°W; 38 m deep), Qassi-sø (64.15°N, -51.31°W; 28 m deep) and Langesø (64.12°N, -51.31°W). Monthly monitoring takes place in Badesø (with fish) and Qassi-sø (without fish).

Monitoring includes ice cover, water chemistry, DOC (dissolved organic carbon), chlorophyll *a*, physical conditions, species composition of plankton, vegetation, bottom organisms, and fish. Physical-chemical parameters, DOC, chlorophyll *a*, phytoplankton and zooplankton are monitored monthly during the ice-free period.

3. Detailed manual

Appendix 1 gives an example of the activities during a monitoring season in Kobbefjord. In the following paragraphs procedures are described in detail for all monitoring elements.

3.1 Plants

3.1.1 The NERO line

The NERO line was established in 2007. It is described in detail in Bay et al. (2008). Surveys of the vegetation transect are done with 5-year intervals. Surveys have been done in 2007 (establishment), 2012 and 2017 and is planned to be done in 2022. The location of the line is shown in figure 3. See

also Bay et al. (2008). The analyses of the transect are only done by trained and experienced botanists.

The NERO line monitoring also includes pin-point analyses as a reference of species composition in the different vegetation types. The sites with pin-point plots are shown in figure 3. Pin-point analyses are performed in seven distinguishable and dominating vegetation types in the Kobbefjord area with five replicas in each vegetation type:

1. Dwarf shrub heath
2. Snow patch
3. Herb slope
4. Fen
5. Copse
6. *Deschampsia* – *Juncus* community
7. Salt marsh




Figure 3. The NERO line. The dots show the positions of the pegs and the colour indicate the vegetation zone northeast of the peg. Numbering of the pegs starts in the south-western corner. The blue dot indicates the position of the characteristic, but rare, plant community dominated by *Deschampsia flexuosa* and *Juncus trifidus*, which is found on south facing, dry slopes. Grey circles mark the sites of pin-point analyses.

Input of data into in-house database at GINR

Data from the Raunkjær circle analyses are entered into a data file named “NERO_line_Raunkjær_data” with the columns: Peg no., Plot no., Year, Month, Day, Observer, Vegetation type, Species name and Raunkjær value. Uncertain species identifications are marked cfr. (confer). Fertile specimens are indicated by (F) added next to the Raunkjær value.

Data from the pin-point analyses are entered into a data file named “NERO_line_pin-point_data” with the columns: Date (YYYY/MM/DD), Vegetation type, Plot (1 – 5), NS (1 – 10), WE (1 – 10), Layer (1 – 3), Species, Status (Live/Dead), Height (cm), Remarks. See NERO line manual (Bay et al. 2008) for further explanation.

Digital pictures, original data sheets, and data files are kept at the Greenland Institute of Natural Resources.

3.1.2 Reproductive phenology

Monitoring of reproductive phenology consists of weekly counts of buds, flowers, and senescent flowers to monitor the proportion of the phenological stages of the species: Northern willow (*Salix glauca* SAL), Trailing azalea (*Loiseleuria procumbens* LOI), and Moss campion (*Silene acaulis* SIL).

Species to be monitored

Three species: *Salix glauca*, *Loiseleuria procumbens*, and *Silene acaulis*. These species have been chosen since they

- Are commonly found in the area.
- Cover a spectrum of different growth forms (deciduous dwarf shrub, evergreen dwarf shrub, and cushion forming herb/dwarf shrub).
- Are comparable to species monitored by BioBasis in Zackenberg.

There are four plots for each species. The size of each plot varies (see table 2) depending on the abundance of individual flowers/catkins of the species in question.

Frequency of sampling

Censuses of *Salix glauca*, *Loiseleuria procumbens*, and *Silene acaulis* are made at weekly intervals in the snow free season (normally May – October). Total counts of flowers are done once a year at peak flowering. Total counts include counting *Eriophorum angustifolium* flowering stems in four plots.

Equipment

- Map with position of study plots (or alternatively GPS)
- Camera
- Data form Appendix 2/ Notebook.

Location and marking of study plots

The location of the 12 plots are shown in Figure 4. The plots are marked with angle pegs in each corner, with one peg (the A peg) carrying a nametag with the plot-ID. They are divided into four subplots (quarters A, B, C, and D) separated by pegs at the centre where the diagonals cross and at the midpoint of each side – see Figure 5. The lettering starts at the corner with the plot-ID and continues clockwise around the centre. Coordinates, dimensions etc. can be found in table 2.




Figure 4. The locations of the plant reproductive phenology plots and plots for annual total counts of flowering shoots for *Salix glauca* (SAL1-SAL4), *Silene acaulis* (SIL1-SIL4), and *Loiseleuria procumbens* (LOI1- LOI4). Note that for *Eriophorum angustifolium* (ERI1-ERI4) only total counts of shoots are monitored. The locations of the *Empetrum nigrum* plots (EMP1-EMP4) for NDVI measurements are included along with the location of the four arthropod plots (marked with grey dots). Coordinates can be found in table 2.

Table 2. Positions and sizes of plant reproductive phenology and *Empetrum nigrum* plots. Positions are given in decimal degrees.

Species	Plot	Latitude	Longitude	Plot dimensions (m)
<i>Eriophorum angustifolium</i>	ERI1	64.1346	-51.3837	4 x 10
	ERI2	64.1312	-51.3873	10 x 10
	ERI3	64.1348	-51.3789	12 x 17
	ERI4	64.1333	-51.3666	5 x 9
<i>Salix glauca</i>	SAL1	64.1325	-51.3729	7 x 11
	SAL2	64.1316	-51.3714	8 x 8
	SAL3	64.1337	-51.3678	6 x 9
	SAL4	64.1374	-51.3741	4 x 5
<i>Silene acaulis</i>	SIL4	64.1361	-51.3681	5 x 12
	SIL1	64.1328	-51.3745	5,5 x 7
	SIL2	64.1364	-51.3703	11 x 11
	SIL3	64.1370	-51.3736	7 x 11
<i>Loiseleuria procumbens</i>	LOI1	64.1323	-51.3759	1.8 x 3.9
	LOI2	64.1316	-51.3705	1.7 x 3
	LOI3	64.1324	-51.3708	1.6 x 2.6
	LOI4	64.1328	-51.3702	1.6 x 3
<i>Empetrum nigrum</i>	EMP1	64.1339	-51.3839	2 x 2
	EMP2	64.1308	-51.3894	4 x 4.1
	EMP3	64.1336	-51.3836	3 x 3
	EMP4	64.1337	-51.3675	2.5 x 2.5




Figure 5. Left: Lettering of subplots in plant phenology plots. The dot indicates the corner with the plot ID. Arrows indicate the clock-wise direction to walk when measuring NDVI. Right: Overview of counting direction applied in phenology subplots A-D. Black dots represent starting points for counting in each subplot and arrows the counting direction. Begin at the peg with the plot-ID and count from left to right, as shown by the arrow. Continue counting by moving back from right to left. This procedure (left -> right, right -> left) is continued until a total of 50 (*Silene*) or 100 (*Salix*, *Loiseleuria*) is reached, or to the end of the subplot, whichever comes first. Then continue to the next subplot, starting at the next peg, indicated by the next black dot in the figure.

Sampling method

The following observations and censuses are entered into the relevant data forms for all plots:

- Time
- Cloud cover
- Plot number
- Snow cover
- Number of buds
- Number of flowers (catkins in *Salix*) – note that for *Salix* both male and female plants can be found in the plots (SAL1, SAL2 and SAL3: both male and female plants; SAL4: only male plants)
- Number of senescent female catkins as indicated by open capsules and/or presence of hairs
- Number of senescent flowers (*Loiseleuria* and *Silene*)
- Annual count of total number of flowers (*Salix*, *Silene* and *Loiseleuria*) and flowering stems (*Eriophorum*)
- Remarks (include information on e.g., larvae or fungi infected leaves)

Before counting, take a photo of the plot in which as much as possible of the plot is included. As a minimum the metal rod with the ID-tag must be included in the photo. Download photos and rename in the following format SpeciesPlot_Date: SSSP_YYMMDD (e.g., ERI1_180628, SAL4_180926).

Data forms are found in Appendix 2. The data from the weekly counts are entered into data files with columns relevant for each species. The basic data are: Year, Month, Day, Photo no., Observer, Species (SAL, LOI, SIL), Plot (1-4), Subplot (A-D), Snow (percent in subplot), Cloud cover (x/8), Buds (actual numbers counted, not percent), Flowers, Senescent (flowers), Total (sum of buds, flowers,

and senescent flowers), and Remarks. Please note, that there are species specific columns in the database files, as *Salix* has separate female and male plants.

During snow melt in May/June, percentage snow cover in each sub subplot is estimated at each sampling trip. If any plant part is visible above the snow layer, the cover is given as 99%. If any ground/vegetation cover is free, no more than 98% can be stated.

When visiting *Silene* plots, samples of a total of 50 flower buds, flowers and/or senescent flowers (or capsules) are recorded within each subplot. In the *Salix* and *Loiseleuria* plots a total of 100 buds, flowers, and senescent flowers (catkins in *Salix*) are recorded. This is done by counting the different phenological stages until a total of 50 (*Silene*) or 100 (*Salix*, *Loiseleuria*) is reached. Start counting at the peg carrying the plot-ID, and count from left to right, as shown in figure 5 (right). The procedure of counting left to right, right to left) is continued until a total of 50 (*Silene*) or 100 (*Salix*, *Loiseleuria*) is reached, or to the end of the subplot, whichever comes first. Continue to the next subplot, starting at the next peg, indicated by the next black dot in figure 5 (right).

In general, flower buds are defined as flowers not yet open while flowers are open giving insects access to the reproductive organs, and senescent flowers as flowers that have lost all petals or with all petals almost or fully faded or brown. In some of the final stages, flower stems from the preceding year may interfere with the counts. However, such old stems are always dry and stiff; stems of this year are soft and fleshy. Handling of plants and flowers must be kept to a minimum to minimize impacts on vegetation. Try to limit disturbance of the plots in general i.e., step as little as possible inside plots, touch as little as possible etc.

For each species, the following sampling procedures apply.

Salix

The sampling unit is catkins, not individual flowers. Most flowers from one catkin emerge the same day, and they also wilt at the same time. Hence, catkins are recorded as male or female buds based knowledge from previous years and/or the presence of old catkins (Figure 6), when no stigmas or anthers are visible. Catkins are recorded as male (Figure 7) and female (Figure 8) flowers as soon as anthers (m) or stigmas (f) are clearly visible/discernible and protruding (they are often both red in the early stages, but the colour may vary). Please notice that the sampling unit may be rooted in a different plot than where they are counted / referred to. If the plant is rooted in e.g., subplot A and has stems with flowers reaching into subplot B, the flowers should be counted in B. Please notice the distribution of male and female plants in the individual plots and subplots in table 3.



Figure 6. *Salix glauca* buds. It is not possible to discriminate between male and female flowers at this stage. Though, based on knowledge from previous years, the buds may be recorded as males or females.



Figure 7. *Salix glauca* male flowers.



Figure 8. *Salix glauca* female flowers.

Both senescent flowers and fruits continue to be recorded as 'flowers' until they have exposed seed hairs (Figure 9) from open capsules. From the time of exposure of the hairs from as much as a single splitting capsule, they are recorded as 'senescent' (note: capsules in "active" catkins may be open but emptied of seed hairs). Notice that fruits may be affected by larvae so that they expose seed hairs from the bottom of the capsules (excreta from the larvae are often visible among the seed hairs, which may also appear denser). These capsules must not be recorded as senescent but should be recorded separately. In Kobbefjord, this was first observed in 2020 (figure 10). Senescent male flowers are not counted.



Figure 9. Senescent *Salix glauca* female flowers with hairs from open capsules



Figure 10. *Salix glauca* female flower eaten by larvae. Orange faecal matter from larvae (marked by arrows).

Fruits infected by fungi (yellow and twisted) should be recorded separately (yet still included in the number for 'flowers', i.e., the infected fruits appear twice in the data forms). This has not been observed in Kobbefjord yet. Also, infections by insects should be recorded.

Table 3. Overview of male and female plants in the *Salix* plots. ^) Female flower observed in 2021. *) Male flowers found in plot until 2016, and not since.

Plot	Subplot	Male plant	Female plant
SAL1	A	✓	✓
	B		✓
	C		✓
	D	✓	✓^
SAL2	A	✓	
	B	✓	
	C		✓
	D		✓
SAL3	A	✓	✓
	B	✓*	✓
	C	✓*	✓
	D	✓	✓
SAL4	A	✓	
	B	✓	
	C	✓	
	D	✓	

Silene

Silene acaulis grows in cushions (Figure 11) and one or a few specimens may dominate the subplots.

Flower buds are reddish or light purple (Figure 12). Senescent flowers (Figure 13) have wilted petals and empty pollen anthers or appear as empty “cups” (Figure 13).

Some years flowers are heavily grazed by insects. In such cases the grazed flowers should be counted as flowers only if parts of the flower are still recognisable (Figure 13). If the remaining parts are not recognisable as e.g., a flower it should not be counted.



Figure 11. *Silene acaulis* cushion.



Figure 12. *Silene acaulis* flowers in the foreground. Buds in different developmental stages are marked with the arrows.



Figure 13. *Silene acaulis* senescent flowers, some with wilted petals (within circle). Buds or flowers eaten by insects in the foreground.

Loiseleuria

Loiseleuria procumbens is a matted shrub with pairs of tiny, oblong, closely set leaves and abundant clusters of small flowers (see Figure 14). In Greenland, plants grow no taller than 10 cm. The plant is creeping, multi-branched, and mat forming, with 2-5 pink, bell-shaped flowers in terminal clusters and evergreen leaves with rolled edges.



Figure 14. *Loiseleuria procumbens* buds (fully closed), flowers (open) and senescent flowers from last year (upper right corner).

Input of data into in-house database at GINR

Data from the weekly censuses are entered into a data file named “Phenology_Plots_YYYY” with the columns: Year, Month, Day, Photo no., Observer, Species (SAL, LOI, SIL), Plot (1-4), Subplot (A-D), Snow (percent in subplot), Cloud cover (x/8), Buds (actual numbers counted, not percent), Flowers, Senescent (flowers), Total (sum of buds, flowers and senescent flowers), Larvae, Fungi and Remarks. Please note that there are species specific columns in the data file.

3.1.3 Total flowering

Species to be monitored

Northern willow (*Salix glauca*), Trailing azalea (*Loiseleuria procumbens*), Moss campion (*Silene acaulis*), and Cotton grass (*Eriophorum angustifolium*). See previous section for descriptions of *Salix glauca*, *Loiseleuria procumbens*, and *Silene acaulis*. *E. angustifolium* is described below.

Eriophorum

The sampling unit is flowering stems, not individual flowers, or spikes. There are two or more spikes on each stem. Figure 15-16 shows different stages of *Eriophorum* flower development.



Figure 15. *Eriophorum angustifolium* with immature spikes (yellow anthers visible).



Figure 16. *Eriophorum angustifolium* with mature spikes (conspicuous white perianth bristles).

Frequency of sampling

Once per season. Total counts of *S. glauca*, *L. procumbens*, and *S. acaulis* are made at peak flowering. The optimal time for total counts of *E. angustifolium* is when most or all spikes have reached senescence.

Equipment

- Map with position of study plots (or alternatively a GPS)
- Pieces of cord totalling 100 m
- Flower sticks
- Tally counters
- Data form Appendix 2/ Notebook.

Location and marking of sampling plots

The plots are divided into four subplots (A, B, C, and D) separated by steel pegs at the centre where the diagonals cross and at the midpoint of each side. The lettering starts at the corner with the plot-ID tag and continues clockwise around the centre. Coordinates, dimensions etc. are found in table 2. The plots are identical with the plant reproductive phenology plots shown in figure 4.

Sampling method

Tighten a cord around each subplot. In large plots, subsections are established by placing two additional cords with about 0.5 or 1 m intervals from one end of each subplot, whereupon the summed number of flower buds, flowers, and senescent flowers, respectively, are counted between each cord. Move one cord at a time and repeat the process until the entire plot is covered. In small plots, sticks may be used instead of cords. In the *Salix* plots, male and female catkins are counted separately. Catkins that have been grazed but can still be identified as either male or female, are included.

Input of data into in-house database at GINR

Data from the yearly registrations are entered into a data file named “Phenology_Plots_YYYY” (in a separate sheet for total counts) with the columns: Year, Month, Day, Photo no., Observer, Species (SAL, LOI, SIL, ERI), Plot (1-4), Subplot (A-D), Snow (percentage in subplot), Cloud cover (x/8), Buds (actual numbers counted, not percent), Flowers, Senescent (flowers), Total (sum of buds, flowers and senescent flowers), and Remarks. Please note, that there are species specific columns in the data file.

3.1.4 Normalised Difference Vegetation Index (NDVI) in plots and along the NERO line

The NDVI-instrument measures the spectral reflectance of the plant canopy. The progression in the vegetation greenness is followed in the plant phenology plots and from 2008-2016 along the vegetation transect (the NERO line). Due to instrument failure (old instrument: handheld Crop Circle TM ACS-210) NDVI along the line has not been measured since. The new instrument (RapidSCAN CS-45) cannot be used for measurements along the NERO line. For information regarding measurements along the NERO line please refer to a previous manual (e.g., Aastrup et al. 2015).

Species or taxonomic groups to be monitored

All plants in the reproductive plant phenology plots and 4 plots with *Empetrum nigrum*.

Frequency of sampling

Plant phenology plots: Weekly in connection with the plant phenology censuses.

Equipment

- Map of plant phenology plots or GPS
- RapidSCAN CS-45
- Camera
- Notebook

Location and marking of sampling plots

Plant phenology plots: See Table 2 and Figure 4.

Sampling method

NDVI is measured using the RapidSCAN CS-45 (See Figure 17). It integrates a data logger, graphic display, GPS, crop sensor and a power source in one small instrument. Please refer to the RapidSCAN

CS-45 manual for in depth information on the instrument. Data is collected and stored on a flash file for later retrieval to a PC using the RapidTALK software.



Figure 17. Equipment for measuring NDVI, the handheld RapidSCAN CS-45. Photo from Holland Scientific homepage.

The procedure is:

1. Turn the RapidSCAN CS-45 on by moving the clicker forwards for a couple of seconds.
2. Wait for GPS signal (may need to turn the instrument on and off a few times).
3. Press "OK" for "Log" mode.
4. Pull clicker backwards once to start logging subplot A. At the end of subplot A, pull clicker backwards to stop logging. Repeat procedure for each subplot following figure 5, left.
5. Press "OK" after logging subplot D.
6. Turn "OFF" between plots.

Scans are conducted by moving the sensor steadily forward (ca. 1 meter per second) approximately 75 cm above the vegetation. Refer to the RapidSCAN CS-45 manual for more information.

The sampling order specified in table 4 must be applied. The RapidSCAN CS-45 does not name the files, so it is extremely important that the sampling order is followed! If for some reason the order is changed, remember to make a clear note in your notebook. Also, always measure all plots in the

order A-D (see Figure 5, left). At each visit, note under Remarks the presence of snow (snow in subplot) and if the vegetation is wet/moist.

Table 4. Walking order of phenology plots including Empetrum plots for measuring NDVI.

Plot abbreviation	Plot name
ERI1	Eriophorum 1
EMP1	Empetrum 1
ERI2	Eriophorum 2
EMP2	Empetrum 2
ERI3	Eriophorum 3
EMP3	Empetrum 3
LOI1	Loiseleuria 1
SIL1	Silene 1
SAL1	Salix 1
SAL2	Salix 2
LOI2	Loiseleuria 2
LOI3	Loiseleuria 3
LOI4	Loiseleuria 4
ERI4	Eriophorum 4
EMP4	Empetrum 4
SAL3	Salix 3
SIL4	Silene 4
SIL2	Silene 2
SIL3	Silene 3
SAL4	Salix 4

All measurements are conducted only on the AB and the CD sides of the plots (Figure 5, left). Place yourself at the plot number plate, just outside the plot. Hold the sensor perpendicular to your walking direction approx. 75 cm above the ground. Switch on the NDVI logger (follow the procedure described above) and walk slowly (approximately 1 m per second) along the sides indicated by arrows on figure 5, left. Use the clicker switch to log the NDVI along each of the four sides in the plot. Only one NDVI value is generated per subplot corresponding to the average NDVI. Hence, four scans (equal to four NDVI measurements) are made in each of the vegetation plots.

Ideally all plots should be measured on the same day. If the vegetation is wet, the measurements must be postponed to the following day.

Input of data into in-house database at GINR

Data is downloaded from the RapidSCAN CS-45 to the computer using the appropriate cable. Each file holds the following variables: Plot (where 1-4 corresponds to the subplot sides A-D), NDRE (Normalized Difference Red Edge Index), NDVI, RE (Reflectance Red Edge), NIR (Reflectance Near Infrared), R (Reflectance Red), Latitude, Longitude, Latitude, Elevation, HDOP (Horizontal Dilution of Precision), Fixtype (always GPS), Date, Time, N (Number of samples per scan), MaxNDRE (Maximum NDRE), MaxNDVI (Maximum NDVI), MinNDRE (Minimum NDRE), MinNDVI (Minimum NDVI), StdNDRE (Standard deviation NDRE), StdNDVI (Standard deviation NDVI), CVNDRE (Coefficient of variation NDRE), CVNDVI (Coefficient of variation NDVI).

Data are entered into a data file named "Plot_NDVI_YYYY" with the following columns: Year, Month, Day, DOY, Date, Observer, Species, Plot, Section, NDRE, NDVI, RE, NIR, R, Latitude, Longitude, Elevation, HDOP, Fixtype, N, MaxNDRE, MaxNDVI, MinNDRE, MinNDVI, StdNDRE, StdNDVI, CVNDRE, CVNDVI.

Pictures are downloaded and renamed according to the procedure described in 3.1.2 Reproductive phenology.

3.1.5 Normalised Difference Vegetation Index (NDVI) in CO₂ flux plots

The progression in the vegetation greenness is followed in the CO₂ flux plots. Spectral reflectance of the plant canopy as well as the incoming radiance is measured using two sensors.

Species or taxonomic groups to be monitored

All plants in the CO₂ flux plots.

Frequency of sampling

Weekly in connection with the CO₂ flux measurements.

Equipment

- Map of CO₂ flux plot positions
- GPS with positions of CO₂ flux plot
- A handheld SpectroSense 2+ system including two light sensors (marked 38291 and 38294 respectively) which calculates the greening index (NDVI). See figure 18
- Data form Appendix 3 or 4 / Notebook.

Location and marking of sampling plots

CO₂ flux plots: See figure 21. The position of the midpoint of the plots is 64.1361°N/51.3831°W.

Sampling method

NDVI in the CO₂ flux plots is measured with the SpectroSense 2+ system using two light sensors. Data is not stored and must be written in the “Remarks” section on either data form 3 or 4.

The procedure is:

1. Fasten the 4 metal legs/rods loosely to the Plexiglas plate using the accompanying screws and bolts. The legs should be fixed but able to turn slightly to fit into the metal frame in the plots – one leg in each corner. This ensures identical measurement height every time.
2. Unscrew the bolt on the light sensors and attach them to the Plexiglas plate as follows: Attach the sensor marked 38294, without a diffuser, facing downwards in the middle of the “square” made out of the 4 bolts from the legs. Attach the sensor marked 38291, with a diffuser, facing upwards outside the “square”.
3. Connect the light sensor marked 38291 (with a diffuser) in the “Current C1/C3” (plug no. 2 from the left) on the device. See figure 18.
4. Connect the light sensor marked 38294 (without a diffuser) in the “Current C2/C4” (plug no. 3 from the left) on the device. See figure 18.
5. Place the 4 legs in the corners of the metal frame in the plot (double check that the sensors are placed correctly). See figure 19.
6. Turn on SpectroSense 2+ with the ON button.
7. Tap the MENU button and scroll down to MENU “6-NDVI” using the arrow keys on the right-hand side. Press ENTER.
8. Initially the NDVI value fluctuates but should reach a fairly stable value after a few seconds. Write the value (with 3 decimals) in data form 3 or 4 in the remarks column.
9. Continue to the next plot. Do not dismantle the setup between plots. The SpectroSense 2+ may turn off automatically at some point. In that case start at #6.
10. When all plots have been measured dismantle the legs from the plexiglas plate and unscrew the light sensors but leave the sensor cables in the device.

The easiest way of sampling the plots is by following the same route as when measuring CO₂ flux. It may be favourable to do the NDVI measuring when the flux measurements are completed.

Input of data into in-house database at GINR

Data is stored in a data file named “ITEX_NDVI_YYYY” with the following columns: Year, Month, Day, Day of Year (DOY), Plot, Treatment, Snow, NDVI and Remarks.

Insert light sensor
marked 38294
here

Insert light sensor
marked 38291
here




Figure 18. The SpectroSense 2+ device for measuring NDVI in the CO₂ flux plots. Attach the light sensors as indicated in the figure.




Figure 19. The light sensors attached to a plexiglas plate with 4 legs positioned in each of the corners of the flux plot metal frame. Notice the light sensor with the diffuser marked 38291 is facing upwards while the sensor without a diffuser (marked 38294) is facing downwards.

3.1.6 CO₂ flux plots

The ratio between the release of CO₂ from respiration and decomposition of organic matter, and the uptake via photosynthesis is measured. The ratio is called Net Ecosystem Exchange (NEE).

Species or taxonomic groups to be monitored

The vegetation in the CO₂ flux plots which is dominated by *Empetrum* heath with *Salix* as subdominant species. The reproductive phenology of *Salix* is followed in all plots. Soil moisture is measured in all plots. Soil temperature is recorded by wireless GeoPrecision mini data loggers.

Frequency of sampling

CO₂ flux is measured weekly. All plots should be measured between 9 AM and 3 PM, and on the same day.

Equipment

- ITEX-chambers (OTCs) incl. bolts (only for set-up)
- Hessian tents (2.2m x 2.2m – only for set-up)
- Electrical tubes for the hessian tents (only for set-up)
- External hexagonal frame for the hessian tents (already located in the six S-plots)
- Wireless GeoPrecision mini data loggers (already located in each plot)
- Wireless GeoPrecision USB-dongle for data retrieval (bring once in autumn)
- EGM-5 – see Figure 22
- TRP-3 probe for measuring CO₂, PAR (photosynthetic active radiation) and temperature
- Handheld computer
- Plexiglass measuring chamber (PMC) – measuring 33x33x34 cm (LxWxH).
- Theta-probe for soil moisture measurements
- Black plastic bag adjusted for the PMC
- Sticky tack
- External 12V battery
- Ruler
- Camera
- Data forms Appendix 3 and 4 / Notebook.

Location and marking of sampling plots

30 plots are located in a mesic dwarf shrub heath dominated by *Empetrum nigrum* and with *Salix glauca* as subdominant species. The heath is facing west.

Figure 20 gives an overview of the site and Figure 21 shows the relative positions of the plots.

The setup consists of 5 treatments each with 6 replicates: Control (C), increased temperature (T, ITEX chambers/OTCs), Shading (S, hessian tents), long growing season (LG, removal of snow during spring) and short growing season (SG, addition of snow during spring).

Temperature is increased in T plots by ITEX chambers (OTCs) measuring 78 cm across the opening and 40 cm in height, see Figure 23. This way temperature is expected to increase 1-2 °C during the growing season (for further information, see Molau & Mølgaard, 1996). The shading treatment (S) is carried out by erecting dome shaped hessian cloth tents over the plot causing approximately 60% reduction of incoming light (Havström et al., 1993).

Short and long growing season plots were intended to be differentiated by respectively adding to and removing snow from SG and LG plots during snow melt in spring causing plots to be exposed

earlier (LG) or later (SG) than in control plots. These two treatments have however not been implemented, but the layout of plots on a slope naturally cause differentiation of snowmelt. The plots thus work as additional control plots.

In each of the 30 plots, a metal frame of 35x35 cm has been inserted permanently into the soil. The frame is used for weekly measurements of CO₂ fluxes in the system by the closed chamber technique. The metal frames were placed in spots where *E. nigrum* and *S. glauca* dominated the vegetation. The metal frames are not to be removed by the end of the season.



Figure 20. Overview of CO₂ flux plot site with Hessian tents for shadowing and ITEX chambers (OTCs) to increase temperature.




Figure 21. Detailed map showing the position of the CO₂ flux plots. The position of the midpoint is 64.1361°N/51.3831°W. For explanation of the abbreviations, please see text.

Sampling method

The CO₂ flux treatments are established as soon as possible in the season (make note of the date). By the beginning of the season, check that the metal frames are level and adjust if needed. Do not adjust the metal frames later or in connection with the CO₂ flux measurements!

The six S-plots are equipped with hessian tents by erecting three electrical tubes in an external hexagonal frame over each plot. Each end of the tubes is placed in the external frame on opposite sides creating a dome. The tubes should be long enough to stand one meter over the vegetation at the highest point of the dome. Cover the plot with a piece of hessian and secure it with tent pegs and rocks. Remember to cover the plot with hessian when doing the light measurement.

The six plexiglas sides of each OTC are bolted together on site. A wireless GeoPrecision mini data logger is already placed approximately 2 cm horizontally into the soil. The data loggers are programmed to log the temperature every 30 minutes.

Before the start of the season, replace the old soda lime (referred to as “CO₂ Scrubber” in the EGM-5 manual) in the EGM-5 with fresh soda lime. This is done in the institute lab. The EGM-5 display provides a “CO₂ scrubber status” (see figure 22) which indicates the status of the soda lime. When the value gets to 20% or less the soda lime must be replaced. For resetting the CO₂ scrubber after soda lime change, please refer to the EGM-5 manual page 38.

Prior to 2021 an EGM-4 instrument was used. Please refer to previous manuals for information on the EGM-4 procedures.

Before beginning the CO₂ flux measurements in a plot take a digital ortho-photo covering the entire area inside the metal frame and take three soil moisture measurements outside the metal frame but inside the plot using the ThetaProbe. See the ThetaProbe manual for further details. Measurements of the frame height is only done three times during the season (beginning, mid and end of season) to avoid unnecessary tear on the vegetation. Frame height is the height (cm) from the upper edge of the metal frame to the ground. Measurements are taken in the middle of each of the four sides with the ruler perpendicular to the metal frame to ensure that the rules don't slide below the soil surface.

Measurement of CO₂ flux

1. The EGM-5 must take a measurement every 10 seconds. This is done by turning on the EGM-5, click the “Main” button, the “Settings” button and choose the “Settings 3 Menu” (done by clicking the “→” in the lower right corner twice), and choose “Host”. The “Interval (sec)” can be changed to 10 by clicking on the black bar. Return to the “Main” page, use the “Back” button and the “←” in the lower left corner).
2. In the field, insert the TRP-3 probe into the plexiglas measuring chamber (PMC). The probe has a cable and a tube with black tape. Connect the cable and the tube from the probe to the EGM-5 (“Probe 1”, and “Gas in”). Then connect the clear tube from the PMC to the EGM-5 (“Gas out”). Seal the entrance to the chamber with sticky tack. Do not turn on the EGM-5 before the probe is connected. Please notice that the sensitivity of the PAR sensor is high resulting in measurements of up to 8 $\mu\text{mol m}^{-2} \text{s}^{-1}$ even though the PMC and PAR sensor is covered with the black casing (the probe as an accuracy of $\pm 10 \mu\text{mol m}^{-2} \text{s}^{-1}$).
3. Connect the computer to the EGM-5 the USB interface.
4. Turn on the EGM-5. The EGM-5 has to heat up to 55°C (it may take 10-15 minutes depending on the surrounding temperature). While heating up, the current internal temperature can be seen on the display as “W tt.t” in the lower mid part of the screen. The “W tt.t” disappears when the internal temperature reaches 55°C. Once the EGM-5 has the correct internal

temperature press “Z” in the upper right corner for the instrument to calibrate. The CO₂ level in the air varies between approximately 380 and 410 ppm. See “Note” below for further on the calibration.

5. Start the program GAS (Gas Analysis Software) programme on the computer. Choose the proper device (EGM-5) and click on the “Connect” button. Choose the available EGM-5 device and click “OK”. The main measurement screen will be displayed.
6. When the EGM-5 is ready for measurements (internal temperature of 55°C and you have made the first calibration) place the PMC in the metal frame in the first plot to be measured. Make sure the PMC handle does not cast a shadow on the TRP-3 probe with the PAR (photosynthetic active radiation) measuring device. While measuring gas flux the EGM-5 automatically measures the PAR and the temperature in the PMC.
7. Press “Start Record” to begin the measurement. 13 measurements are needed in both light and dark conditions. The counting to 13 is not done automatically by the instrument, so you will have to follow the screen regularly.
8. Press “Save Log” to save the file in the designated folder. Remember to name your file correctly i.e., name of the plot (e.g., 1c, 2sg etc.) including information on light conditions e.g. 1c_d (dark measurement). The two measurements (light and dark) are saved in separate files.
9. Between the light and dark measurement, the PMC must be carefully aerated to make sure the CO₂ level returns to that prior to measuring. Monitor the CO₂ concentration on the EGM-5 or the computer. Place the PMC on the frame again and cover it completely with the black plastic bag. Repeat #7 and #8 to make the second measurement in the plot.
10. When the light and dark measurements in the plot have been completed lift the PMC and aerate it while walking to the next plot.
11. When all plots have been measured turn off the EGM-54 and computer.
12. Make sure the hessian-tents (and ITEX chambers) are securely fastened before leaving the site.

NOTE: The EGM-5 is set to calibrate (performing “zeros”) manually. It is thus important that you press the “Z” in the upper right corner of the touch display on the EGM-5 once in a while. This could be after each plot measurement, or at least after 5 plots before going to a new plot group.

Measurement of soil moisture

Take three measurements in each plot by pressing the sensor into the soil, read out the soil moisture (press “read” for measurement), and enter data into the form Appendix 3. Press “esc” before next measurement.

Reproductive phenology of Salix

Reproductive phenology of *Salix glauca* is followed in all plots according to the procedure described in section 3.1. Please remember to distinguish between flowers inside and outside the metal frame permanently inserted in the soil. Data are entered into the form Appendix 4.

Species composition of plots

Species composition of the vegetation inside the metal frame is monitored every 5 years, coinciding with the survey of the NERO line. This is done by pin-point analyses and supplementary registration of all remaining species (those not hit by pin-point intercept) within the metal frame.

Laboratory work

None.

Input of data into in-house database at GINR

Data is saved directly on the handheld computer. The raw files are saved in a separate folder (named by the date of measurement) and renamed to include the type of treatment including information on dark measurement status e.g., 1c.txt and 1c d.txt (where the “d” is referring to a dark measurement). Data is stored in a data file named “C_flux_YYYY” and reorganized and supplemented with the following columns: Year, Month, Day, DOY, , Date, Observer, Plot, Treatment, Light, Photo_no, Cloud cover, Time, PLOT NO, RECORD NO, CO₂, AIRPRESSURE, FLOW RATE, H₂O, H₂O TEMP, O₂, SYSTEM ERROR, AUX VOLTAGE, PAR, TSOIL, TAIR, RH, P1, P2, P3, P4, P5, Soil_moisture_1-3, Frame height, and Remarks. All files are merged into one, covering the whole season.




Figure 22. Top: The back of the EGM-5 instrument used for measuring CO₂ concentrations. Bottom, left: The touch display at the top of the EGM-5. Bottom, right: The TRP-3 temperature/PAR sensor.

Download digital pictures and rename them to include plot name and date (e.g., ITEX_1C_090602). Save under “Photos” in the folder named “C-flux”.

Data from the wireless GeoPrecision mini data logger is downloaded once a year at the end of the season. Use the USB-dongle connected to the handheld computer to download the data files from the loggers. Save the individual files using the plot name (e.g., 5C) and save in a separate folder named “C-flux_temperature”. If a connection cannot be made to a logger due to code failure, the general code for access is “GLORIA”.




Figure 23. Measurement of CO₂ flux in an ITEX-hexagon open-top chamber with a measuring chamber (PMC) fitted with a TRP-2 probe connected to the EGM4. This setup resembles the “old” instrument and probe, though the setup with the new EGM5 and TRP-3 probe is similar. The black plastic bag is used for dark (respiration) measurements.

3.1.7 UV-B exclusion

The impact of ambient UV-B radiation on the vegetation is studied in a mesic dwarf shrub heath by placing filters approximately 10 cm above the vegetation. At the time of peak plant growth, the chlorophyll *a* fluorescence is measured as an indicator of plant health. The impact of UV-B exclusion has not been measured since 2017. In 2018, the experiment was set up, but no measurements were done.

Species or taxonomic groups to be monitored

A mesic dwarf shrub heath (facing WSW) dominated by crowberry (*Empetrum nigrum*) and with dwarf birch (*Betula nana*) and bog bilberry (*Vaccinium uliginosum*) as subdominant species. The effect of UV-B radiation is measured on *Betula nana* and *Vaccinium uliginosum*.

Frequency of sampling

Measurements of chlorophyll fluorescence of leaves of *Betula nana* and *Vaccinium uliginosum* is carried out three times with one-week intervals at the peak of plant growth (week 29 to 31).

Equipment

- HandyPea fluorimeter (PEA: Photosynthesis Efficiency Analyzer) – See figure 24 and further description below.
- 80 leaf clips

- Data form Appendix 5 / Notebook
- Camera
- 5 Frames with UV-B filter (Mylar film, 0.25 mm - with exclusion of UV-B)
- 5 Frames of filter control (Teflon film - without exclusion of UV-B).



Figure 24. Handy PEA with clips.

The Handy PEA chlorophyll fluorimeter consists of a control unit. The chlorophyll fluorescence signal received by the sensor head during recording is digitized within the Handy PEA control unit. Up to 1000 recordings may be saved in the memory of Handy PEA chlorophyll fluorimeter. Saved data can be viewed onscreen but has to be transferred to a computer for storage and further analysis.

The sensor unit consists of an array of 3 ultra-bright red LED's optically filtered to a peak wavelength of 650 nm, which is readily absorbed by the chloroplasts of the leaf. The LED's are focused via lenses onto the leaf surface to provide even illumination over the area of leaf exposed by the leaf clip (4 mm diameter).

Location and marking of sampling plots

The UV-B plots are located west of the CO₂ plots. Figure 25 and 26 gives overviews of the plots.

There are three series of plots with five replicates:

1. Control - no treatment: C1-C5
2. UV-B filter (Mylar film, with exclusion of UV-B): B1-B5
3. Filter control (Teflon film, without exclusion of UV-B): F1-F5.

Each treatment plot measures 60 cm x 60 cm; the plots are marked with aluminium tubes at each corner and covered with a frame with the appropriate filter placed approximately 10 cm above the

vegetation. During summer, the vegetation may grow as tall as the filter, which should then be lifted.



Figure 25. Overview of UV-B plots.

Sampling method

Before establishing the UV-B plots, filters on frames are checked carefully, and changed if necessary. There are two filter types, Teflon (filter control: the thinnest and most flexible film) and Mylar (excludes UV-B: thicker and less flexible). Frame positions are given by small sticks within each plot. UV-B plots are checked regularly during the entire field season and repaired if necessary. Specifically, filters in the UV-B plots must be checked after heavy rain or wind.

By the end of the field season, all equipment at UV-B plots is taken down and brought back to the institute.




Figure 26. Schematic presentation of the location of UV-B plots.

Sampling procedure

1. Select five green, “healthy-looking” leaves, or shoot-tips of *Betula nana* and *Vaccinium uliginosum* in each plot.

2. Mount leaf clips on all *Betula nana* leaves – preferably without removing the leaf from the branch. Mount on one species at a time. Make sure that the leaf is visible through the hole in the clip and push the shutter to cover the hole, so the leaf material is in complete darkness.
3. Keep the shutters closed for at least 30 minutes. The closure time may be longer.
4. Switch on the Handy PEA, open “Main menu” and turn the arrow on the screen to “Measure”.
5. Fit the sensor head to the clip; uncover the hole by pushing the shutter back. Start measuring by pushing “OK” on the Handy PEA or push the black button on the sensor head. During the measurement, a number of parameters appear on the screen. Note that the “Fv/Fb” should be about 0.8. If something goes wrong step three must be repeated before carrying out a new measurement.
6. Accept to store the measurement. Note which measurement number corresponds to each plot.
7. Repeat the sampling now measuring on *Vaccinium uliginosum* leaves.
8. Take a photo of each plot at each measurement round.

Input of data into in-house database at GINR

When all measurements have been completed, data must be transferred to a computer by use of the Handy PEA programme. Make sure that all data has been transferred to the computer before clearing the memory in the Handy PEA.

Data are downloaded from the Handy PEA using the PEA Plus software, and the raw files are saved in a separate folder named to include the date of the measurements (e.g., yymmdd.pcs). Data are stored in a data file named “HandyPea_YYYY” and are supplemented with the following columns: Year, Month, Day, Observer, Species, Treatment, and File no.

3.2 Arthropods

Surface living arthropods are captured in yellow pitfall traps. Flying arthropods are captured in window traps.

Species to be monitored

All taxonomic groups of arthropods.

Frequency of sampling

The traps are emptied weekly. If bad weather prohibits visits to the fjord or proper handling of the samples, the traps must be emptied on the earliest day of convenience.

Equipment

For field work:

- Map of the plots (or alternatively a GPS)
- 16 yellow (Pantone no. 108U) plastic cups (pitfalls), 10 cm in diameter, 8 cm deep (permanent position in plot). At the beginning of the season check if all cups are placed properly
- Extra yellow pitfalls for emptying the permanent pitfalls
- 4 window traps (permanent locations)
- A garden trowel with sharp edge (most relevant in the beginning of the season)
- 5 L container for water
- 5 L container for wastewater

- Detergent: Odour free detergent (Coop Änglemark Bluecare Dishwashing fluid, concentrated, without perfume, colour and preservation agent)
- Salt (NaCl) without iodine and anti-caking agent
- 20 Metal pegs (to be used in the fen area)
- 1 lady's stocking per emptying bout (light brown/skin coloured, 20 denier)
- A pair of flat tweezers
- 20 plastic 100 ml containers with lids
- 2 L of 70% ethanol (1400 ml 96% ethanol + 600 ml demineralised water)
- Squeeze bottle with tip (for rinsing the stocking and pitfall with alcohol)
- Disposable syringes for removal of surplus water
- Ethanol resistant labels (see Appendix 14)
- Data form Appendix 6 /notebook.

Location and marking of sampling plots

The location of the study plots are shown on Figure 27. Each plot measures 10 x 20 m and is made up of eight 5 x 5 m squares marked with metal pegs in each corner. Each plot is identified with an ID-label, and subplots (with one trap each) are lettered A-H clockwise from the number plate, see figure 28. Starting from the 2021-season, only 4 pitfall traps (A-D) per plot are opened.

Sampling set up

A set of four pitfall traps has been established in each plot (figure 28). Each trap is composed of two plastic pit-fall cups fitting into each other, so that the upper one can be lifted and emptied without disturbing the surrounding soil. The traps have been permanently positioned within squares, each of 5 x 5 m. Previously (prior to 2016) the traps were moved randomly each year. This procedure has been abandoned due to the slow rate at which the vegetation and soil surface regenerate.

Repositioning of the traps each year caused considerable damage to the plots and would eventually render them non-representative of the surrounding areas.

One cup per pitfall is placed upside-down during the winter. At the start of the season (i.e., on the day when the traps have appeared from the snow), new clean (washed with a little biodegradable, non-perfumed detergent) upper cups are placed inside the 'wintering' ones.

If there is any risk of the cups floating due to water in the lower cup, two metal pegs must be placed along each cup to keep them in position.

The upper cup of the trap is filled $\frac{2}{3}$ - $\frac{3}{4}$ with water (approx. $\frac{1}{2}$ litre needed per station) added three drops of detergent and a spoonful of salt as killing agent, preservation and to prevent freezing.

Two window traps are set up at both Art2 and Art3. They are set up to catch flying insects in all compass directions (N-S and E-W). The traps are fixed on the ground with two angle irons permanently inserted in the soil. The traps are filled ca. $\frac{3}{4}$ with water and added 2 spoonsful of salt and 5 drops of detergent.




Figure 27. Location of arthropod plots. Art1: *Empetrum nigrum* heath, Art2: *Fen*, Art3: *Betula nana* / *Salix glauca* heath, Art4: *Abrasion*. Window traps are located at Art2 and Art3. Phenology plots are marked with grey dots for reference.

Emptying the traps

Prepare and print labels on water resistant paper before field work (Appendix 14). Labels should contain date, plot (Art1-4 or Wart1-4) and subplot (A-D). Remember to write the date on the label if that was not included in the printing. Samples from each of the pitfall traps and window traps are kept separate.

1. Open a 100 ml container and insert the correct pre-printed label.
2. Stretch the stocking over the top of a spare cup.
3. Pour the trap liquid through the stocking into the spare cup.
4. Check the pitfall for remaining arthropods. Should any remain in the trap, flush with ethanol down into the 100 ml container.
5. Reposition the pitfall in the soil.
6. Empty the catch from the stocking into the 100 ml container by turning the stocking upside down on top of the container.
7. Rinse the inverted stocking with ethanol using a small squeeze bottle.
8. Check for arthropods in the stocking. All remaining arthropods must be removed carefully from the stocking using tweezers and put into the container.
9. Fill the container approx. $\frac{3}{4}$ with 70% ethanol. More is needed if there are many insects. In extreme cases, the sample may be divided between 2 containers.
10. Pour the water (now with no arthropods) back into the pitfall repositioned in the soil.

Window traps are emptied using a plastic spoon with a mesh net. Before emptying a window trap a spare cup is filled (approx. 50%) with water from the trap. Use the spoon with the net to catch all arthropods in the trap. Rinse the spoon in the spare cup now and then. The spare cup is emptied as described above (1. – 10.) when all arthropods have been fished out of the window trap.

Water may need to be added (to compensate for evaporated water), removed (after rain) or replaced (e.g., if the water smells or if foxes have urinated or defecated in the pitfall). When water is replaced make sure to add a little salt and detergent. Bring all contaminated water back in a wastewater container and pour it into the sink in the “lab-room” in the main cabin.

Bring extra pit-fall cups on each round, along with equipment for setting up traps, in case a trap has been destroyed, e.g., by a fox. Any failures such as flooded or floating cups, fox faeces etc. must be recorded. This includes occurrence of fungi and/or algae in the water. In that case, cup and water must be replaced. Cups contaminated by e.g., faeces, fungi, or algae can be cleaned with detergent and ethanol in the lab and reused once clean.




Figure 28. Schematic diagram showing positions of arthropod subplots A-H. Starting from the 2021-season, only 4 pitfall traps (A-D) per plot are opened.

Note the full hour of the day when the traps in each plot are emptied.

During snow melt (and after snow fall in late summer / autumn) the snow cover (%) is estimated for each subplot. If any plant part is visible above the snow layer, the cover is given as 99%. If any ground/vegetation cover is free, no more than 98% can be stated.

Never touch the traps with mosquito repellent or sun cream on your fingers!

Ending the season

At the end of the catching season the trap liquid must be collected from all the traps. The wastewater is emptied in the sink in the “lab-room” in the main cabin (or brought back to the institute if the drainage system has been closed for the winter). Leave the angle irons for the window traps in the soil. Arthropod samples are kept at GINR before they are shipped to the Department of Bioscience, Aarhus University, Silkeborg, for sorting and identification.

Laboratory work

None.

Input of data into in-house database at GINR

After the weekly emptying of the pit-fall traps, the following data are entered into a data file named “Arthropods_YYYY”: Year, Month, Day, DOY, Hour, Observer, Trap (Art or WArt), Plot (1-4), subplot (A-D), Snow, Days (days since last emptying), Remarks. Under Remarks, date of opening and closing together with relevant observations about the traps are stated. This includes any disturbance that may influence the efficiency of the traps such as flooding, drying out, icing, dirt, faeces, adding/removing water, and vandalism by mammals including humans.

After sorting, the total number of individuals per group is entered into a data file (please refer to the GEM database metadata file for further information).

3.3 Microarthropods

Species to be monitored

All microarthropods.

Frequency of sampling

Sampling of microarthropods is done in 8 plots (4 habitats with 2 replicates; figure 29). Four samples from each plot are collected three times during the season corresponding to: spring (after snowmelt), summer and autumn (before the snow appears). Extraction is very slow in wet samples. Therefore, samples should only be taken when soil moisture is low i.e., a few days after a rain event.




Figure 29. Location of plots for microarthropod sampling.




Figure 30. Microarthropods sampling grid in 10x10 m plots with grid size: 0.5 x 0.5 m.

Equipment

- Map/GPS with positions of plots
- Soil auger
- 34 microcosm tubes made of Plexiglas (height 5.5 cm and diameter 6 cm)
- 64 DBIdut lids (size 89B)
- Knife to cut roots etc.
- Transportation box
- Table with random sampling points (Appendix 15)

The yearly sampling programme consists of collecting microarthropod samples from:

4 habitats * 2 plots * 4 subsamples * 3 sampling occasions = 96 samples.

Each plot is marked with 4 pegs. To ensure enough undisturbed sampling points for several years each plot is divided into a ½-meter square grid with two axes (x, y; figure 30). The peg with the ID-tag (MArt1-8) is the starting corner for the square grid (x, y = 0m, 0m). The table with random sampling points provides 4 sets of random numbers (x, y coordinates) from where the soil samples are to be taken. The same set of random numbers are used for all 8 plots on each sampling occasion.

Sampling method

1. Two microcosm tubes are inserted into the soil auger, which is then closed and ready for use.
2. The point of sampling is found using the random sampling table.
3. The soil auger is placed vertically at the sampling point, so it touches the soil surface.

4. At sites with dense vegetation, it may be necessary to use a knife to cut around the soil auger before pushing it down into the soil or peat. Take care not to damage and compress the soil/peat core.
5. Push the soil auger vertically 5.5 cm downwards so that the lowest tube is just filled with soil. The soil surface has to level the upper rim of the lowest tube. The soil auger is open in the top so that you can follow how the soil is pressed into the tube. The function of the upper tube is only to keep the lower tube fixed. While pushing the soil auger down, turn it from side to side thereby avoiding compression of the soil in the tube.
6. Tilt the soil auger from side to side to loosen the soil core at the bottom and carefully pull the soil auger including the soil core up.
7. Open the soil auger and carefully remove the tube including the soil core. Place a labelled DBIdut lid at the top immediately to keep organisms on the soil from escaping. Make sure that the lid is properly marked with the sample name (e.g., MArt4C).
8. Turn the tube around and cut surplus soil away with the knife so the soil surface levels the bottom of the lower tube. Place a DBIdut lid in the bottom of the tube.
9. Place the tubes in a box with the top of the sample facing upwards.

Store the samples at low temperature in a shaded place and avoid excess bumping during transportation. On arrival to the laboratory, the samples are stored in the dark at 5°C until extraction no later than two days after sampling.

Laboratory work

Extraction of microarthropods

The extractor holds 36 samples, so all samples from one sampling batch are extracted at the same time. When the extractor heats the samples from above the microarthropods will move towards colder soils lower in the samples. They will eventually move through the mesh and drop into the benzoic acid.

Equipment

- Extractor with temperature sensor and data logger
- Manual for extractor
- Insulation foam
- Cooling circulator with anti-freeze fluid and tubes (for connection to extractor)
- 32 soil samples
- 32 meshes with a mesh size of 1x1 mm
- 32 extraction cups with saturated benzoic acid (14.5 g benzoic acid and approximately 1 ml detergent per 5 L)
- Detergent
- 32 lids for extraction cups (only to be used if for some reason you are not able to pour the samples into the 100 ml containers the same day the extraction is finished)
- 96% ethanol (may be denatured if pure ethanol is not available)
- 100 ml containers with lids
- Ethanol resistant labels (see Appendix 15)

Extraction procedure

1. One day before extraction: Start the cooling circulator connected to the extractor as the samples should not be extracted at bottom temperatures higher than 5°C. Fill the base of

- the extractor with water (will be cooled by the circulating anti-freeze fluid running in the pipes at the base of the extractor).
2. At the day of extraction: Pour 0.5 cm of a saturated solution of benzoic acid and detergent (14.5 g benzoic acid in 5 L water added roughly 1 ml detergent) in each of the extraction cups. Mark each extraction cup with the sample name.
 3. Put the extraction cups into the extractor.
 4. For each sample: Take a microcosmos tube containing a soil sample. Carefully remove the upper lid and place the mesh on the tube with the sample.
 5. Place a suitable cup above the soil sample unit and turn the cup with the sample up-side-down.
 6. Remove the DBIdut lid from the bottom (now at the top after turning the microcosmos tube and mesh) and sweep surplus soil down into the cup.
 7. Place the microcosm tube with the mesh on the corresponding extraction cup in the extractor. Make sure the topsoil surface is facing downwards.
 8. Pour any surplus soil into the soil sample.
 9. Place the insulating material around the samples when all samples are in place in the extractor. The insulation around the tubes must be placed carefully so that no soil particles will drop into the extraction cups. Fill surplus holes in the insulation with the black insulation foam circles.
 10. Connect one temperature sensor into the soil of one of the samples in the extractor for regulation of temperature.
 11. Close the extractor.
 12. Turn on the extractor and press the green start button. The extractor will now heat the samples according to this schedule:
 - 30°C for 48 hours
 - 40°C for 48 hours
 - 50°C for 48 hours
 - 60°C for 24 hours, terminated manually by switching off the power supply, but it may be continued until all the samples are dry on the down-facing surface on the mesh. The cooling system should ensure that the temperature of the benzoic acid solution is minimum 4°C and maximum 20°C throughout the extraction.
 13. The extraction is stopped manually by turning the power off.
 14. Check that the samples are dry on the surface facing downwards after termination of the pre-programmed extraction process. If some samples are still wet, continue the extraction at 60°C until the samples are dry. Samples with high organic content can be divided in two, to ensure complete extraction.
 15. Turn off the extractor and cooling circulator.
 16. Throw the soil away.
 17. Empty the benzoic acid holding the microarthropods from the extraction cups into a 100ml container. Clean the sides of the cup with 96% ethanol using a squeeze bottle, spraying the sides down to ensure any remaining microarthropods are flushed into the container.
 18. Make sure that the amount of ethanol added is in the proportion one part benzoic acid to two parts of ethanol (resulting in approximately 70% ethanol). If necessary to obtain this proportion divide the sample into two plastic cups. Put a pre-printed water-resistant label into the container with the date and name of the sample.
 19. Brush the nets clean. Wash the tubes.
 20. Store the samples with lids closed tightly until identification at Department of Bioscience, Aarhus University, Silkeborg.

3.4 Birds

Monitoring of birds consists of two elements: Breeding phenology of small passerines on an ad hoc basis and weekly samplings of bird observations at permanent points.

3.4.1 Breeding phenology of passerines

Species to be monitored

The passerine bird species Northern wheatear (*Oenanthe oenanthe*), Snow bunting (*Plectrophenax nivalis*), Lapland Bunting (*Carduelis flammea*), and Common Redpoll (*Calcarius lapponicus*) are monitored in the study area indicated in figure 2 and from census points as shown in figure 31.

Frequency of sampling

During June and July on an ad hoc basis. Nests of breeding passerines are located ad hoc, and the located nests are followed as frequently as possible until the chicks have left the nest.

Equipment

- Binoculars (10x)
- GPS
- Data form Appendix 7 / Notebook
- Sticks for marking of nests

Sampling method

At all visits at located nests note:

- Date
- Species
- Number of eggs/chicks
- GPS position
- Take close-up photo of the nest and chicks.

Input of data into in-house database at GINR

The position of nests is entered into a data file named “Bird_nests_YYYY” with the following columns: Species, Date, Observer, GPS-position, Number of eggs, number of chicks, and Remarks.

3.4.2 Point sampling

The primary objective of this study is to monitor the birds in the Kobbefjord valley. It is, however, also a very good opportunity to watch for other kinds of wildlife. The focus is on the small passerines.

Species to be monitored

The passerine bird species Northern wheatear (*Oenanthe oenanthe*), Snow bunting (*Plectrophenax nivalis*), Lapland Bunting (*Carduelis flammea*), and Common Redpoll (*Calcarius lapponicus*).

Frequency of sampling

Weekly during the entire field season.

Equipment

- Binoculars (10x)
- Clock
- Data form Appendix 8 / Notebook.

Location and marking of sampling plots

The observation points are located by GPS (Figure 30 and table 5).

Table 5. Positions of bird observation points A-M. Positions are given in decimal degrees.

Point	Latitude	Longitude
Bird A	64.134685	-51.385105
Bird B	64.135155	-51.391187
Bird C	64.134592	-51.396234
Bird D	64.132390	-51.393590
Bird E	64.131052	-51.389160
Bird F	64.129385	-51.378330
Bird G	64.131761	-51.379398
Bird H	64.132669	-51.374116
Bird I	64.134509	-51.363874
Bird J	64.135639	-51.355553
Bird K	64.133636	-51.344558
Bird L	64.132841	-51.336278
Bird M	64.131031	-51.326204




Figure 31. Bird observation points. The Kobbefjord Research Station is indicated by the orange symbol close to the observation point Bird A.

Sampling method

Record species (one of the four passerine species), number, sex (male, female, unknown) and age (juvenile, adult, unknown), special behaviour, etc. At each point sit for 5 minutes before starting the 5 minutes observation period. If other species are encountered, they must be recorded on the “Random observations” list (Appendix 9). The list is normally found on the board in the main cabin.

Input of data into in-house database at GINR

The number of birds observed at each observation point is entered into an data file named "Birdcountings_YYYY" with the following columns: Year, Month, Day, DOY, Observer, Time, Point (A-M), Obsperiod (2), Cloud cover (x/8), Species (NW, SB, LB, or RP), Age (Juvenile, Adult, Unknown), Gender (M, F, Unknown), Number, and Comments.

3.5 Mammals

Species to be monitored

All mammal species.

Frequency of sampling

Ad hoc continuously during entire field season.

Equipment

- Binoculars (10x)
- Notebook.

Location and marking of study plot

Entire monitoring area, see Figure 2.

Sampling method

Keep watch for everything with fur. Record geographical position, group size, sex and age, special behaviour etc. All personnel are encouraged to supply observations and submit these on the observation sheet in the main cabin.

See observation sheet in Appendix 9 (Random observations).

Input of data into in-house database at GINR

Data from Appendix 9 are entered into a data file named "Random Obs_YYYY." with the following columns: Year, Month, Day, Observer, Location, GPS-position, Species, and Remarks.

3.6 Lakes

Physical-chemical and biological (phytoplankton, zooplankton, fish and macrophytes) parameters.

Parameters monitored monthly

- Ice cover
- Water temperature
- Water transparency (Secchi depth)
- pH
- Conductivity
- Chlorophyll *a*
- Total nitrogen
- Dissolved nitrogen (nitrite/nitrate and ammonium)
- Total phosphorus
- Dissolved phosphorus (phosphate)
- Dissolved organic carbon (DOC)
- Phytoplankton (all taxonomic groups)
- Zooplankton (all taxonomic groups)

3.6.1 Water samples

Frequency of water sampling

Five times per year, during the ice-free period (normally June – October). First sampling must be done as early in the season as possible – preferably, when there is still some ice on the lake. Sampling is done approximately every 4 weeks. If the ice-free period is shorter than 5 months, the period between samplings is reduced to fit five samplings during the ice-free period.

Ice cover is recorded continuously via automatic cameras operated by GeoBasis.

Location and marking of sampling spots

The two sampling lakes are located in the Kobbefjord catchment area in the inner part of Kobbefjord (Badesø / Kangerluarsunnguup Tasia: 64.13°N, 51.36°W, 38 m deep, and Qassi-sø: 64.15°N, 51.31°W, 28 m deep; Figure 32). There is one sampling station at the deepest position in each of the lakes. The station is marked with a buoy anchored at the bottom. At the buoy-line 2 temperature loggers are mounted at 2m and 10m depth, respectively. If possible, a sediment trap is attached at the buoy-line at 10m (Qassi-sø) and 15m (Badesø). It has to be collected and stored during winter. Positions of the sampling stations are saved in the GPS.




Figure 32. Location of the sampling station in each of the two monitored lakes.

Equipment for monthly sampling

- A rubber dinghy with oars and outboard motor
- Pump for the dinghy
- Plastic bottles for sediment (two per lake, only during the last visit – and only if sediment trap is active)
- A water sampler
- 30 litre tub

- Zooplankton filter (20 µm mesh)
- Plastic funnel for 60 ml zooplankton bottles
- Squeeze bottle
- 2 x 60 ml brown glass bottles including 2% Lugol solution for zooplankton (2.5 ml Lugol) and phytoplankton (1 ml Lugol) samples
- Plastic bottles for water chemistry (250 ml), DOC (250 ml) and chlorophyll *a* (2 l)
- Tinfoil
- Secchi disc with rope with metre measure
- HOBO Optic USB Base Station and TidbiT v2 Coupler for the HOBO (only needed in the lab)
- HOBO TidbiT Temperature Logger (only in the beginning of the season)
- Depth sounder with thermometer
- Life jacket
- Survival suit
- Data forms Appendix 10 and 11.

Equipment for laboratory work

- Whatmann GF/C filters (47 mm) for Chl *a*
- Tinfoil
- Small plastic tubes for Chl *a* filters
- Equipment for analysing DOC
- Detailed list of chemicals etc. to be used for analysis – see Appendix 13.

Sampling methods

At each sampling date, time, cloud cover (x/8), wind speed (calm, light, weak, hard, etc.) and ice cover (% of lake area) is recorded. Use data form Appendix 10 for recording data in the field.

Bring water sampler, tub, Secchi disc, depth sounder, sampling bottles, oars, and a filled outboard. Go to the sampling station and tie the dinghy to the buoy.

Before sampling, rinse all sampling gear and bottles with lake water. Measure the transparency with the Secchi disc at an accuracy of 0.1 m on the light side of the boat. Keep the face close to the water surface; lower the disc until it disappears and pull it slowly upwards until it is just visible – this is the Secchi depth.

Take a depth integrated pooled water sample of approximately 25 litres from 0.5 m below the surface to approximately 0.5 m above the bottom. Start from the top and avoid any sediment in the sample. If the lake sediment is disturbed and occur in the sampler a new sample must be taken. Wait 10 minutes or move the boat away from the first sampling spot. Measure the water temperature both at the surface and at every water sample through the water column.

While in the dinghy, stir the water in the tub well and take:

1. Approx. 100 ml water sample in a 250 ml bottle for water chemistry
2. Approx. 150 ml water sample in a *new/unused* 250 ml bottle for DOC (dissolved organic carbon)
3. 2 litres of water for chlorophyll *a* measurements
4. One 60 ml sample for phytoplankton enumeration (preserved with 1 ml Lugol solution)

5. 15 litres filtered through 20 µm filter. Release the filter and transfer the collected zooplankton to a 60 ml bottle (preserved with 2.5 ml Lugol solution. Rinse well with water from the pooled sample).

All sample bottles are kept dark (e.g., the samples to be used for chlorophyll *a*, water chemistry and DOC measurements should be wrapped in tinfoil) and cold.

At last visit in the season: Empty the sediment traps (see text below) into two plastic bottles, and place in correct position in water column again (traps have not been active in the lakes for several years). At the same time, change the two TidbiT v2 temperature loggers with two empty loggers. Data is downloaded in the lab. Make sure the temperature loggers are logging with 1-hour intervals. One logger is fitted 2 m below the surface and named "Lake name_2m". The other one is placed 10 m below the surface, named "Lake name_10m". Every 4 years the TidbiT v2 loggers are replaced (two new sets have been set up in 2020).

After each sampling, the dinghy is placed at the same spot at Qassi-sø together with the outboard and other sampling gear. The dinghy is thoroughly secured to the big rock using rope. At Badesø the dinghy is secured after each sampling and taken back to the storage cabin after the last visit in the season. The outboard motor used at Badesø is taken back to the storage cabin after each sampling. The outboard at Qassi-sø is brought back at the end of the season.

Laboratory work

Before filtering for chlorophyll *a* (Chl *a*), absolute conductivity (accuracy 1 µS), temperature (°C) and pH (accuracy 0.1 pH unit) is measured. This is done in the cabin. For chlorophyll measurements 2 x 1 litre from each lake is filtered (exact volume in ml is recorded) through two 47 mm GF/C filters. The filters are folded, placed in separate test tubes, extracted in 10 ml 96% ethanol, wrapped in tinfoil, and analysed according to Dansk Standard, DS2201 (Appendix 13).

Dissolved organic carbon (DOC)

In the laboratory, cover your "workspace" / table with tinfoil to ensure a carbon-free environment. Prepare 2 test tubes ("burned" 6 hours at 450 °C in Muffle oven, lids prewashed in 0.1 M HCl) labelled with lake ID, date and containing 150 µl 2 M HCl. Use tweezers to place a 25 mm (also "burned" 6 hours at 450°C in Muffle oven) Whatman GF/F filter in a filter bracket. A 60 ml plastic syringe is mounted with a 15 cm teflon tube. The syringe is filled with 60 ml of the water sample, which is then used to rinse the syringe i.e., flush the water through the syringe by pressing the piston and letting the water out in the sink. Afterwards the syringe is filled with min. 50 ml of water sample. The teflon tube is removed and the filter bracket is mounted on the syringe. Press the sample through the filter (for flushing of filter) until 40 ml is left. The test tubes are each filled with 15 ml of filtered sample (filtered through the same filter) and the lids are tightened.

The test tubes are wrapped in tinfoil and stored in the fridge until analysis. The filter is removed from the filter bracket and thrown away. The syringe, teflon tube and the filter bracket are rinsed in ELGA water and air dried. DOC is analysed at Department of Bioscience, Aarhus University, Silkeborg.

Water samples for water chemistry are frozen until analyses at Department of Bioscience, Aarhus University, Silkeborg.

Zooplankton and phytoplankton samples are stored in darkness – not frozen. These samples are sent to Department of Bioscience, Aarhus University, Silkeborg following the last sampling.

3.6.2 Submerged vegetation monitored annually

The annual monitoring of the submerged vegetation takes place in August. The method to be used for the macrophyte studies is a transect investigation to obtain a relatively good overall description of submerged macrophyte distribution, density, and diversity. The macrophyte diversity is very low and only 2-3 species/taxa are found in the monitored lakes: *Callitriche hamulata* (intermediate water-starwort), moss, and in 2018 *Myriophyllum alterniflorum* (alternate water-milfoil) was found in Badesø close to the river outlet from Langesø. In 2020 a single plant of *Hippuris vulgaris* (mare's tail) was found close to the beach on the southwest side of Badesø.

In each lake, approx. 150 observation points are included. If great spatial variability in plant cover occurs, more points may be needed to provide an adequate description.

The observation points should follow straight lines from one shore towards the other (parallel to each other and perpendicular to the shore) to a depth of 2x maximum depth of the growing vegetation (min. 5 meters and max. 15 meters). All observation points are marked on a GPS.

At each observation point, water depth, total macrophyte coverage (index between 0 and 5, see next paragraph), and the species/taxa are recorded (Appendix 12).

The investigation is undertaken by two people; one conducts all observations (degree of total coverage, species identification, and water depth). The other person operates the dinghy, enters all observations in the data sheet and saves the coordinates of all observation points into a GPS. Observations are made using a plant rake with 1 catch at each observation point and estimating the coverage using a 0-5 scale, where 0 = 0 %; 1 = >0 - 5 %; 2 = >5 - 25 %; 3 = >25 - 50 %; 4 = >50 - 75 % and 5 = >75 - 100 %.

Example of basic analyses of data from the transect investigation

The following observations were obtained in a transect investigation of a 75 ha lake: 25*0%, 32*3%, 40*15%, 23*38%, 19*63% and 11*88%, i.e. a total of 150 observations.

The relative plant-covered area (RPA) is calculated as follows:

$(25*0\% + 32*3\% + 40*15\% + 23*38\% + 19*63\% + 11*88\% / 150 = 25\%),$

This requires that all observations have been made at the same equidistant intervals all over the lake. If relatively more observations have been made in dense near-shore macrophytes, the observations should be weighted before RPA is calculated.

Equipment for annual vegetation sampling

- Plant rake with robe
- Handheld depth sounder
- GPS
- Data form Appendix 12.

3.6.3 Parameters monitored every 5th year

- Fish (taxonomic groups, population and for isotopes)
- Macroinvertebrates (taxonomic groups, abundance and for isotopes)
- Phytoplankton and zooplankton for isotopes
- Sediment (paleo-analyses of chironomids, diatoms and cladocerans).

Fish

In each lake a maximum of 9 (1.5 m deep) sinking Lundgren biological multi mesh gill nets are deployed for approximately 16-18 hours (figure 33). Gill nets (3 in each zone) are set in the littoral (close to the bottom at 2-2.5 m depth), in the pelagic (in the middle of the water column at 15-20 m depth) and at the bottom (benthic nets at 18-20 m) in the late afternoon. All nets are set up parallel to the shore using anchors (one at each end) and buoys (one at each end for littoral and benthic nets and 2 additional buoys along the net for the pelagic nets). Nets are gathered the following morning. The catch is treated per net and per net type (littoral, pelagic, benthic). When removing fish from more nets from the same depth it is vital that they are kept in separate buckets until they are processed. And further, if nets from different depths are sorted it becomes even more crucial to separate them.

Each fish is given a consecutive number, identified to species (Arctic char *Salvelinus alpinus* or Three-spined stickleback *Gasterosteus aculeatus*) and sex, length and weight is recorded.

- Tissue samples (dorsal muscle) for isotope analyses: For Arctic char, samples are taken from approximately 20 fish per size class (≤ 20 cm and >20 cm). The tissue samples are frozen in plastic vials and marked with lake name, date, fish number (same as above) and content. Three-spined sticklebacks are sampled as whole fish.
- The stomach is sampled from Arctic char. They are preserved in 96 % ethanol in vials or 100, 200 or 300 ml jars, depending on the size of the stomach. The container is marked with lake name, date, fish number (same as above) and content.
- Otoliths are sampled from Arctic char. These are kept in paper envelopes and marked with lake name, date, fish number (same as above) and content.




Figure 33. Lundgren gill net set up with anchors at the bottom and buoys at the surface.

Sediment/paleo samples

Four sediment cores are sampled in each of the two lakes using the kajak sampler. One from a depth of approx. 15m (in part of the lake with more or less low inclination/horizontal bottom) and three close to the shore (taken by hand). Be careful not to disturb the sediment cores when unscrewing the core from the sampler. When the cores are taken on board there must be a sharp threshold between the sediment and the water phase. If this is not the case it indicates that the core may have tipped over at the bottom and the sample must be re-taken.

On shore, surface sediments (0-1 cm) from the three near-shore cores are pooled in a 250 ml plastic jar and marked with lake name, date, number of cores and depth (0-1 cm). Surface sediment from the core sampled at 15m is kept separate but marked as above. It has not yet been possible to collect the sediment samples from the larger depth due to the surface of the lake bottom. Thus, we only sample the sediment closer to the shore.

Samples are kept cool and dark.

Samples are analysed at Department of Bioscience, Aarhus University, Silkeborg.

Isotope sampling and analyses

- a. Samples of fish (see above)
- b. Benthic invertebrates: Pelagic and littoral invertebrates are kept separate. Surface sediment is sampled with a sweep net (littoral and in macrophytes) or an Ockelmann-sledge. Samples are rinsed as much as possible in the dinghy before being put into a bucket (several samples from one habitat can be pooled in one bucket as a qualitative sample). On shore, the samples are filtered through a 212 µm mesh and invertebrates are sorted into groups in small glass jars (~5 ml) and marked with lake name, date.
- c. Zooplankton is sampled as two fractions (> 140 µm and > 500 µm). Both nets are pulled after the dinghy until enough material has been collected. Samples are put into 20 ml vials.
- d. Phytoplankton: A large amount of water is filtered through first a 80 µm mesh followed by a 11 µm mesh and samples are put into 20 ml vials (this has not been done in Kobbefjord yet as it requires a mechanical pump to filter the large amount of water – several hundred litres).
- e. Benthic algae: The top 5 mm from one sediment core is put into a vial. If animals are present, they should be removed.
- f. Stones: Fist large stones are collected in the littoral zone and epiphytes are scraped into a vial.
- g. Periphyton: Plants are selected randomly. Plants are washed in a 1 litre jar and periphyton is scraped off. The water including the periphyton is filtered on a 20 µm mesh. Animals are removed and the filtrate is put into a vial.
- h. Macrophytes: A sample is taken of the dominating taxa. Periphyton is removed and roots are avoided. The sample is put into a vial.

All samples are kept frozen in small glass jars. Keep enough material for 3 replicates of all samples.

Laboratory work

Identifying and counting of benthic invertebrates

Samples are shipped to Department of Bioscience, Aarhus University, Silkeborg, together with the plankton samples. The samples are sorted, identified, and counted to species or genus level.

Isotope sample

All isotope samples (fish, invertebrates, plankton, periphytes, macrophytes) are prepared for analyses, i.e. each sample is freeze dried, homogenized and a 5 mg sub sample is weighed in a tin capsule (ready for analyses). Preparation for analyses is undertaken in Nuuk. The isotope samples are analysed at the AU Department of Bioscience Stable Isotope Analytic Center.

Equipment for every 5th year field work (done in 2008, 2013 and 2018)

The equipment is stored in 4 large boxes at GINR. The contents are split into: Fishing gear, Equipment to bring to the field, Lab equipment and “Spare box”. Make sure everything is carefully packed down and replenished when necessary. The gillnets must only be stored when they are completely dry.

Fish

- Gillnets
- Floaters and robe
- Sinks and robe
- Poles for nets
- Buckets for fish
- Field schemes
- Ruler, balance
- Gear for the isotope sampling (see below)

Sediment

- Kajak sampler
- Sediment cores and rubber stoppers
- Equipment to process sediment cores

Isotopes

- Tweezers and scalpels
- Rubber gloves
- Ethanol and squeeze bottles
- Phytoplankton net (prefer 11 µm net but 20 µm is OK)
- Zooplankton net, 140 µm
- Vertical net for large zooplankton, 500 µm
- Sweep net for invertebrates, 500 µm
- Sorting trays
- Plastic vials for invertebrates
- Coolers

3.7 Random observations of birds and mammals

Species to be monitored

All bird species and mammals (Arctic foxes, caribou, and hares).

Frequency of sampling

Continuously during entire field season.

Equipment

- Binoculars
- Spotting scope
- Notebook
- Entire monitoring area
- Data form Appendix 9.

Sampling method

Keep watch for everything with feathers and fur. Record flock size, sex and age, special behaviour, geographical position etc.

3.8 Disturbance

Parameters to be monitored:

‘Person-days’ spent in the area, aircraft activity over the area, boat trips to and from the area, discharges (burning of waste, human discharges into the fiord). All of these can only be monitored during periods with personnel at the station.

4 Storage of data

4.1 Data at GINR

Data collected during the season are downloaded/saved at the Greenland Institute of Natural Resources when returning to Nuuk. All data are typed into or transferred to the specific Excel files immediately. Data is stored in a [Teams folder](#). Each year has a corresponding folder (e.g., “BioBasis_Nuuk_2022”) where each sub-programme has a folder of their own, containing their data files, where the collected data is entered. At the end of the season all data is sent to the Department of Bioscience, Aarhus University, Roskilde for storage (see 4.2 Data for the GEM database as different layout and formats may be used). All original data files are stored on the servers at GINR. Furthermore, written material is stored at GINR along with collected specimens until processed at the appropriate facilities in Denmark.




Figure 34. Layout of the in-house database at GINR. Each BioBasis subprogramme has its own folder. The root folder also contains files with other types of information e.g., a field calendar, field journal and the printed forms to be used.

4.2 Data for the GEM database

Deadline for data delivery is 1 May. Before delivery, all data are quality checked and reformatted for the GEM database. Dates are given as YYYY-MM-DD and the file format is to be a text document (.txt). A table of all files to be sent is shown in table 6. Column and element descriptions can be found in the meta-data files “Biobasis_Nuuk_Column_description” and “Biobasis_Nuuk_Element_description” in the [GEM Database Teams folder](#). Please be aware that for some of the data files the layout has to be changed substantially to fit the form required by the GEM database manager.

All of the datasets in the GEM-database have their own unique doi’s (digital object identifier).

Data can be downloaded from <https://data.g-e-m.dk/>. [To download data, you will have to register / set up an account.](#)

Table 6. Files for the GEM database.

Data	Database name
Arthropod	BioBasis_Nuuk_Arthropd
Bird census	BioBasis_Nuuk_Bird_Census_points
C-flux measurements	BioBasis_Nuuk_CFlux
Lakes, temperature	BioBasis_Nuuk_Lakes_Temperature
Lakes, chemistry	BioBasis_Nuuk_Lakes_PhysChem
Lakes, vegetation	BioBasis_Nuuk_Lakes_Vegetation
Lakes, phytoplankton	BioBasis_Nuuk_Lakes_Phytoplankton
Lakes, zooplankton	BioBasis_Nuuk_Lakes_Zooplankton
UVB measurements (not since 2016)	BioBasis_Nuuk_Leaf_Flourescence
Microarthropod	BioBasis_Nuuk_Microarthropod
NDVI in C-flux plots	BioBasis_Nuuk_NDVI_CFlux
NDVI in phenology plots (until 2016)	BioBasis_Nuuk_NDVI_PhenologyPlots
NDVI in phenology plots (from 2018)	BioBasis_Nuuk_NDVI_PhenologyPlots_New
Phenology in C-flux plots	BioBasis_Nuuk_Phenology_CFlux
Phenology in Loiseleuria plots	BioBasis_Nuuk_PhenologyPlots_Loiseleuria
Phenology in Salix plots	BioBasis_Nuuk_PhenologyPlots_Salix
Phenology in Silene plots	BioBasis_Nuuk_PhenologyPlots_Silene
Phenology – total counts in LOI, SAL, SIL, and ERI plots	BioBasis_Nuuk_PhenologyPlots_TotalCount
PinPoint analyses in C-flux plots	BioBasis_Nuuk_PinPoint_CFlux
Raunkjaer analyses on NERO-line	BioBasis_Nuuk_Raunkjaer_NERO_line

5 References

Bay C, Aastrup P, Nymand J. 2008. The NERO line. A vegetation transect in Kobbefjord, West Greenland. National Environmental Research Institute, Aarhus University, Denmark. 40p. NERI Technical Report no. 693. <http://www2.dmu.dk/Pub/FR693.pdf>.

Fredskild B, Mogensen G. 1997. ZERO line. Final Report 1997. A description of the plant communities along the ZERO line from Young Sund to the top of Aucellabjerg and the common plant communities in the Zackenberg valley, Northeast Greenland. Greenland Botanical Survey & Botanical Museum, University of Copenhagen, 36p.
[http://www.zackenberg.dk/fileadmin/Resources/DMU/GEM/Zackenberg/pdf/zero-line-final-report .pdf](http://www.zackenberg.dk/fileadmin/Resources/DMU/GEM/Zackenberg/pdf/zero-line-final-report.pdf)

Havström M, Callaghan TV, Jonasson S. 1993. Differential growth responses of *Cassiope tetragona*, an arctic dwarf-shrub, to environmental perturbations among three contrasting high- and subarctic sites. *Oikos* 66:389-402.

Molau U, Mølgaard P. 1996. ITEX Manual. 2nd Edition. Danish Polar Center. 85p.

Jansen MAK, Gaba V, Greenberg BM. 1998. Higher plants and UV-B radiation: balancing damage, repair and acclimation. *Trends in Plant Science* 3:131-135.

Post E, Forchhammer MC, Bret-Harte S et al. 2009. Ecological dynamics across the Arctic associated with recent climate change. *Science* 325:1355-1358.

6 List of scientific and technical consultants

General matters

Katrine Raundrup, GINR, Manager of BioBasis Nuuk (+299 361228, kara@natur.gl)

Maia Olsen, GINR (+299 361282, maol@natur.gl)

Ida Bomholt Dyrholm Jacobsen, GINR (+299 361232, idja@natur.gl)

Arthropods

Zdenek Gavor, Elin Jørgensen, Aarhus University (+45 87158838, zg@bios.au.dk)

C-flux calculations

Mikhail Mastepanov, Aarhus University (+46 739360967, mikhail.mastepanov@bios.au.dk)

Freshwater ecology

Torben Lauridsen, Aarhus University (+45 87158762, tll@bios.au.dk)

Microarthropods

Paul Henning Krogh, Aarhus University (+45 87158831, phk@bios.au.dk)

Vascular plants

Ida Bomholt Dyrholm Jacobsen, GINR (+299 361232, idja@natur.gl)

7 Appendix

Appendix 1 Example of annual sampling protocol

Date	May	June	July	August	September	October
1					Phen; Art; K2 vegetationstransek	
2	Phen; Art; NDVI; Birds			NERO; Phen; Art; NDVI		
3	opsætning ART, Phen, NDVI			UVB		
4			Birds; NERO Phen; Art; NDVI			C-Flux; Phen; Art; NDVI; ½ NERO
5			Besøg MASTERCLASS	C-Flux;		
6		DRYAS; Tea Time; Mart				Birds
7			Besøg MASTERCLASS		C-Flux; Phen; Art; NDVI; Birds	
8		C-flux			Amerikanske stud; MART	
9		NERO; Phen; ½ Art; NDVI				
10		Birds; ½ Art				
11	C-Flux; Phen; NDVI; DRYAS; ½ NERO					C-Flux; Art pakkes sammen
12	Art; opsætning Wart; UV b			Art;		K1
13			C-Flux; Phen; Art; NDVI	C-Flux; Phen; NDVI		
14		Birds; DRYAS	Kristian kommer		Birds; NERO; Art 3 + 4; NDVI;	
15		Sirlus	Birds; Totaltælling		NDVI (er1-emp2); ART 1+2; Jordprøver	
16		C-Flux; Phen; Art; NDVI		C-Flux; Birds;	Jordprøver	
17	Birds; Phen; NDVI			Phen; Art; NDVI		
18						
19	Art			NERO		c-flux
20		Birds; DRYAS	C-Flux; Mart; Birds; NERO		C-Flux; Phen; Art; NDVI	c-flux
21				GU dag i Kobbefjord	Kamera Nordlandet	
22			Phen; Art; NDVI; UVB			
23	K1; Phen; NDVI; NERO; ITEX kh+NDVI			C-Flux; Phen; Art;		send prøver til Dk og Cflux Temp data
24		NERO; Phen; Art; NDVI; K1				
25				NERO; NDVI; Veg. trans ½ Badesø; K1		
26			Phen; NDVI; K1+K2	K2, ½ vegtrans Badesø	Birds;	
27	C-Flux; ; Art; Birds		C-flux; Art	Muslinger		
28		K2	C-flux; Birds; UVB; zooplankton;	Birds - GU i Kobbefjord	Phen; Art; NDVI, K2	
29		Phen; totaltælling; ART		C-flux; Kommunal bestyrelse		
30		Birds; C-flux; NDVI				
31			UVB			
32						

Appendix 2 Data form for plant phenology monitoring

Date:				Observer:								Page 1/2	
Plot	Time	Photo	Cloud cover	#	Snow	Buds	Male flower	Female flower	Hair/ Scene.	Fungus	Larvae	Total	Remarks
ERI1				A									
				B									
				C									
				D									
EMP1				A									
				B									
				C									
				D									
ERI2				A									
				B									
				C									
				D									
EMP2				A									
				B									
				C									
				D									
ERI3				A									
				B									
				C									
				D									
EMP3				A									
				B									
				C									
				D									
LOI1				A									
				B									
				C									
				D									
SIL1				A									
				B									
				C									
				D									
SAL1				A									
				B									
				C									
				D									
SAL2				A									
				B									
				C									
				D									

Date:				Observer:				Page 2/2					
Plot	Time	Photo	Cloud cover	#	Snow	Buds	Male flower	Female flower	Hair/ Scene.	Fungus	Larvae	Total	Remarks
LOI2				A									
				B									
				C									
				D									
LOI3				A									
				B									
				C									
				D									
LOI4				A									
				B									
				C									
				D									
ERI4				A									
				B									
				C									
				D									
EMP4				A									
				B									
				C									
				D									
SAL3				A									
				B									
				C									
				D									
SIL4				A									
				B									
				C									
				D									
SIL2				A									
				B									
				C									
				D									
SIL3				A									
				B									
				C									
				D									
SAL4				A									
				B									
				C									
				D									

Appendix 3 Data form for CO₂-flux monitoring

BioBasis Nuuk 2021			Date:		EGM-4: A/B					Observer:				
Time	Plot	Record nos.		Cloud %/8	Photo	Soil_moisture			Frame_height				Remarks	
		Light	Dark			1	2	3	1	2	3	4		
:	1C	-	-											1C
:	1SG	-	-											1SG
:	1S	-	-											1S
:	1LG	-	-											1LG
:	1T	-	-											1T
:	2SG	-	-											2SG
:	2LG	-	-											2LG
:	2S	-	-											2S
:	2C	-	-											2C
:	2T	-	-											2T
:	3C	-	-											3C
:	3T	-	-											3T
:	3SG	-	-											3SG
:	3LG	-	-											3LG
:	3S	-	-											3S
:	4LG	-	-											4LG
:	4SG	-	-											4SG
:	4S	-	-											4S
:	4T	-	-											4T
:	4C	-	-											4C
:	6S	-	-											6S
:	6SG	-	-											6SG
:	6C	-	-											6C
:	6T	-	-											6T
:	6LG	-	-											6LG
:	5T	-	-											5T
:	5S	-	-											5S
:	5LG	-	-											5LG
:	5SG	-	-											5SG
:	5C	-	-											5C

Appendix 4 Data form for plant phenology in CO₂-flux plots

Salix phenology in C-flux plots

BioBasis Nuuk 2021

Observer:

Date:

Inside										Outside										Plot
Plot no	Snow	Buds	M flowers	F flowers	Hairs	Fungus	Larvae	Buds	M flowers	F flowers	Hairs	Fungus	Larvae	Total	NDVI	Remarks				
1C																	1C			
1SG																	1SG			
1S																	1S			
1LG																	1LG			
1T																	1T			
2SG																	2SG			
2LG																	2LG			
2S																	2S			
2C																	2C			
2T																	2T			
3C																	3C			
3T																	3T			
3SG																	3SG			
3LG																	3LG			
3S																	3S			
4LG																	4LG			
4SG																	4SG			
4S																	4S			
4T																	4T			
4C																	4C			
6S																	6S			
6SG																	6SG			
6C																	6C			
6T																	6T			
6LG																	6LG			
5T																	5T			
5S																	5S			
5LG																	5LG			
5SG																	5SG			
5C																	5C			

Appendix 5 Data form for UV-B effects monitoring

BioBasis Nuuk 2021 Date: _____ Observer: _____

Betula				Vaccinium			
Block	Treatment	Rec. no.	Remarks (DOH)	Block	Treatment	Rec. no.	Remarks (DOH)
1	B (Mylar)	-		1	B (Mylar)	-	
1	C (kontrol)	-		1	C (kontrol)	-	
1	F (film)	-		1	F (film)	-	
2	B (Mylar)	-		2	B (Mylar)	-	
2	C (kontrol)	-		2	C (kontrol)	-	
2	F (film)	-		2	F (film)	-	
3	B (Mylar)	-		3	B (Mylar)	-	
3	C (kontrol)	-		3	C (kontrol)	-	
3	F (film)	-		3	F (film)	-	
4	B (Mylar)	-		4	B (Mylar)	-	
4	C (kontrol)	-		4	C (kontrol)	-	
4	F (film)	-		4	F (film)	-	
5	B (Mylar)	-		5	B (Mylar)	-	
5	C (kontrol)	-		5	C (kontrol)	-	
5	F (film)	-		5	F (film)	-	

Appendix 6 Data form for arthropod monitoring

BioBasis Nuuk 2021				Date:		Observer:		Windtraps					
Time	Plot no.	Sample	Snow	Pitfall opened	Remarks								
:	1	A											
		B											
		C											
		D											
:	2	A				WArt1	NS						
		B				WArt2	EW						
		C											
		D											
:	3	A				WArt3	NS						
		B				WArt4	EW						
		C											
		D											
:	4	A											
		B											
		C											
		D											

Appendix 8 Data form for passerine bird monitoring

BioBasis Nuuk 2021				Date:			Observer:								
Time	Site	Cloud X/8	Snow bunting			Lapland bunting			Red poll			Northern wheatear			Remarks / Other birds
			M	F	UK	M	F	UK	M	F	UK	M	F	UK	
:	A														
:	B														
:	C														
:	D														
:	E														
:	F														
:	G														
:	H														
:	I														
:	J														
:	K														
:	L														
:	M														

When arriving at a new site, sit for 5 min before starting your observations.

(if you see e.g. a fox, hare, caribou, peregrine falcon, white-tailed eagle - incl. tracks and feces - please make a note here)

[illegible]

Appendix 10 Data form for lake monitoring (Badesø)

BioBasis Nuuk 2021

Lake: Badesø

Observer:

Date	Time	Ice cover	Cloud cover	Wind speed	Secchi depth	Depth
	:					

Temp. surface	pH	Conductivity/temp C

Dybde (m)/Temp (°C)
0.5 /
4 /
8 /
12 /
16 /
20 /
25 /
31 /
/

Sample date:

Filter date:

Analysis date:

Analysis by:

Chl a measurements

L=

H=

Sample 1

Before HCl=

ml filtered:

After HCl=

Sample 2

Before HCl=

ml filtered:

After HCl=

RINSE BOTTLES WELL WITH LAKE WATER BEFORE USE

App. 25 litres of pooled water from 0.5m below the surface to 0.5m above the bottom

Samples taken from pooled water to:

Water chemistry (ca. 100 ml. water in 250 ml bottle)

Chl a (2 litres, wrap in tin foil)

Phytoplankton (60ml preserved with 1 ml lugol)

Zooplankton (15 litre filtered through 20µm filter into 60ml bottle with 2.5ml lugol)

DOC (fill half a 250ml bottle and wrap with tin foil)

pH and conductivity are measured on the pooled water

Page 67 of 73

Appendix 11 Data form for lake monitoring (Qassi-sø)

BioBasis Nuuk 2021		Lake: Qassi-sø			Observer:	
Date	Time	Ice cover	Cloud cover	Wind speed	Secchi depth	Depth
	:					

Temp. surface	pH	Conductivity/temp C

Dybde (m)/Temp (°C)
0.5 /
3 /
6 /
9 /
12 /
16 /
20 /
24 /
/

Sample date:	
Filter date:	
Analysis date:	
Analysis by:	
Chl a measurements	
L=	H=
Sample 1	ml filtered:
Before HCl=	After HCl=
Sample 2	ml filtered:
Before HCl=	After HCl=

RINSE BOTTLES WELL WITH LAKE WATER BEFORE USE

App. 25 litres of pooled water from 0.5m below the surface to 0.5m above the bottom

Samples taken from pooled water to:

- Chl a (2 litres, wrap in tin foil)
- DOC (fill half a 250ml bottle and wrap with tin foil)
- Water chemistry (ca. 100 ml. water in 250 ml bottle)
- Phytoplankton (60ml preserved with 1 ml lugol)
- Zooplankton (15 litre filtered through 20µm filter into 60ml bottle with 2.5ml lugol)
- pH and conductivity are measured on the pooled water

Appendix 12 Data form for submerged vegetation monitoring in lakes

Veg_Lake: _____ Dato _____ Side 1/

Waypoint	Dybde	Vandstjerne	Mos	Bemærkninger
1				
2				
3				
4				
5				
6				
7				
8				
9				
10				
11				
12				
13				
14				
15				
16				
17				
18				
19				
20				
21				
22				
23				
24				
25				

Veg_Lake: _____ Dato _____ Side 2/

Waypoint	Dybde	Vandstjerne	Mos	Bemærkninger
26				
27				
28				
29				
30				
31				
32				
33				
34				
35				
36				
37				
38				
39				
40				
41				
42				
43				
44				
45				
46				
47				
48				
49				
50				

Appendix 13 Measuring chlorophyll *a* in the laboratory

Remember that chlorophyll is sensitive to light, thus filtering and measurements must take place in dimmed lighting.

1. Method: fluorometric measurement with ethanol extract

Filtering and extraction

- A known volume of water sample (preferably 2 x 1 litres) is vacuum filtered using GF/C filters. The pressure is not to exceed 0.3 bars or phytoplankton might be destroyed. Filtering time should not exceed ½ h. If so, a smaller volume is to be filtered. Flush the filtering funnel in ELGA-water at the end of filtration.
- Fold the filter and insert in test tube with tweezers.
- Ad 10 ml of ethanol and seal with a lid.
- Shake the sample gently (large air bubbles should be avoided) and extract for 18-24 hours in darkness. If the samples cannot be analysed the following day, they may be frozen after addition of ethanol. After thawing the samples are extracted in darkness for 18-24 hours. Do not leave the samples in the freezer for longer periods of time since the chlorophyll will gradually break down.

Measurements using a fluorometer

- Turn the fluorometer on
- Press the filter to the bottom of the test tube using a tweezer
- Spin samples in the centrifuge at 2000 rpm for 10 minutes
- Carefully transfer the samples to a cuvette (ca. 2 cm from the upper edge) and measure in the fluorometer
- Ad 200 µl (3 drops) of 1 M HCl and turn the cuvette upside down a few times
- Measure again after 2 minutes

2. Reagents

Ethanol, 96% (v/v) density = 0,808

Hydrochloric acid, HCl, 1M

3. Waster and cleaning up

Samples can be poured into the sink when flushing with tap water. The plastic test tubes with filter paper are thrown in the regular dustbin. The 2 litre bottles, filtering funnels, tweezers and cuvettes are rinsed 3 times in ELGA-water.

4. Reference

Modified from Dansk Standard, DS2201 using a fluorometer instead of a spectrophotometer

Appendix 14 Labels for arthropod samples

W GREENLAND Kobbefjord, BioBasis Date: 20/08 2021 ART1A GreenlandInstNatRes AarhusUni, DeptBioSci	W GREENLAND Kobbefjord, BioBasis Date: 20/08 2021 ART2A GreenlandInstNatRes AarhusUni, DeptBioSci	W GREENLAND Kobbefjord, BioBasis Date: 20/08 2021 ART3A GreenlandInstNatRes AarhusUni, DeptBioSci	W GREENLAND Kobbefjord, BioBasis Date: 20/08 2021 ART4A GreenlandInstNatRes AarhusUni, DeptBioSci
W GREENLAND Kobbefjord, BioBasis Date: 20/08 2021 ART1B GreenlandInstNatRes AarhusUni, DeptBioSci	W GREENLAND Kobbefjord, BioBasis Date: 20/08 2021 ART2B GreenlandInstNatRes AarhusUni, DeptBioSci	W GREENLAND Kobbefjord, BioBasis Date: 20/08 2021 ART3B GreenlandInstNatRes AarhusUni, DeptBioSci	W GREENLAND Kobbefjord, BioBasis Date: 20/08 2021 ART4B GreenlandInstNatRes AarhusUni, DeptBioSci
W GREENLAND Kobbefjord, BioBasis Date: 20/08 2021 ART1C GreenlandInstNatRes AarhusUni, DeptBioSci	W GREENLAND Kobbefjord, BioBasis Date: 20/08 2021 ART2C GreenlandInstNatRes AarhusUni, DeptBioSci	W GREENLAND Kobbefjord, BioBasis Date: 20/08 2021 ART3C GreenlandInstNatRes AarhusUni, DeptBioSci	W GREENLAND Kobbefjord, BioBasis Date: 20/08 2021 ART4C GreenlandInstNatRes AarhusUni, DeptBioSci
W GREENLAND Kobbefjord, BioBasis Date: 20/08 2021 ART1D GreenlandInstNatRes AarhusUni, DeptBioSci	W GREENLAND Kobbefjord, BioBasis Date: 20/08 2021 ART2D GreenlandInstNatRes AarhusUni, DeptBioSci	W GREENLAND Kobbefjord, BioBasis Date: 20/08 2021 ART3D GreenlandInstNatRes AarhusUni, DeptBioSci	W GREENLAND Kobbefjord, BioBasis Date: 20/08 2021 ART4D GreenlandInstNatRes AarhusUni, DeptBioSci
	W GREENLAND Kobbefjord, BioBasis Date: 20/08 2021 WART1 GreenlandInstNatRes AarhusUni, DeptBioSci	W GREENLAND Kobbefjord, BioBasis Date: 20/08 2021 WART3 GreenlandInstNatRes AarhusUni, DeptBioSci	
	W GREENLAND Kobbefjord, BioBasis Date: 20/08 2021 WART2 GreenlandInstNatRes AarhusUni, DeptBioSci	W GREENLAND Kobbefjord, BioBasis Date: 20/08 2021 WART4 GreenlandInstNatRes AarhusUni, DeptBioSci	

Appendix 15 Labels for microarthropod samples

W GREENLAND
Kobbefjord, BioBasis
Date: ___/___/2021
MARTIA
GreenlandInstNatRes
AarhusUni, DeptBioSci

W GREENLAND
Kobbefjord, BioBasIs
Date: __/__/2021
MART3A
GreenlandInstNatRes
AarhusUni, DeptBioSci

W GREENLAND
Kobbefjord, BioBasis
Date: ___/___/2021
MARTSA
GreenlandInstNatRes
AarhusUni, DeptBioSci

W GREENLAND
Kobbefjord, BioBasis
Date: ___/___/2021
MART7A
GreenlandInstNatRes
AarhusUni, DeptBioSci

W GREENLAND
Kobbefjord, BioBasis
Date: ___/___/2021
MART1B
GreenlandInstNatRes
AarhusUni, DeptBioSci

W GREENLAND
Kobbefjord, BioBasis
Date: ___/___/2021
MART3B
GreenlandInstNatRes
AarhusUniv, DeptBioSci

W GREENLAND
Kobbefjord, BioBasis
Date: ___/___/2021
MARTSB
GreenlandInstNatRes
AarhusUnl, DeptBioSci

W GREENLAND
Kobbefjord, BioBasis
Date: ___/___/2021
MART7 B
GreenlandInstNatRes
AarhusUni, DeptBioSci

W GREENLAND
Kobbefjord, BioBasis
Date: ___/___/2021
MARTIC
GreenlandInstNatRes
AarhusUniv, DeptBioSci

W GREENLAND
Kobbefjord, BioBasIs
Date: ___/___/2021
MART3C
GreenlandInstNatRes
AarhusUnl. DeptBioSci

W GREENLAND
Kobbefjord, BioBasis
Date: ___/___/2021
MARTSC
GreenlandInstNatRes
AarhusUnl. Dept BioSci

W GREENLAND
Kobbefjord, BioBasis
Date: ___/___/2021
MART7C
GreenlandInstNatRes
AarhusUni, DeptBioSci

W GREENLAND
Kobbefjord, BioBasis
Date: ___/___/2021
MARTID
GreenlandInstNatRes
AarhusUni, DeptBioSci

W GREENLAND
Kobbefjord, BioBasIs
Date: ___/___/2021
MARTSD
GreenlandInstNatRes
AarhusUnl. DeptBioSci

W GREENLAND
Kobbefjord, BioBasis
Date: ___/___/2021
MARTSD
GreenlandInstNatRes
AarhusUnl. DeptBioSci

W GREENLAND
Kobbefjord, BioBasis
Date: ___/___/2021
MART7D
GreenlandInstNatRes
AarhusUni, DeptBioSci

W GREENLAND
Kobbefjord, BioBasis
Date: ___/___/2021
MARTZA
GreenlandInstNatRes
AarhusUni, DeptBioSci

W GREENLAND
Kobbefjord, BioBasIs
Date: ___/___/2021
MART4A
GreenlandInstNatRes
AarhusUni, DeptBioSci

W GREENLAND
Kobbefjord, BioBasis
Date: ___/___/2021
MARTSA
GreenlandInstNatRes
AarhusUni, DeptBioSci

W GREENLAND
Kobbefjord, BioBasis
Date: ___/___/2021
MARTBA
GreenlandInstNatRes
AarhusUni, DeptBioSci

W GREENLAND
Kobbefjord, BioBasis
Date: ___/___/2021
MART2B
GreenlandInstNatRes
AarhusUni, DeptBioSci

W GREENLAND
Kobbefjord, BioBasis
Date: ___/___/2021
MART4B
GreenlandInstNatRes
AarhusUniv, DeptBioSci

W GREENLAND
Kobbefjord, BioBasis
Date: ___/___/2021
MARTSB
GreenlandInstNat Res
AarhusUnl. Dept BioSci

W GREENLAND
Kobbefjord, BioBasis
Date: ___/___/2021
MARTB
GreenlandInstNatRes
AarhusUni. Dept BioSci

W GREENLAND
Kobbefjord, BioBasis
Date: ___/___/2021
MART2C
GreenlandInstNatRes
AarhusUni, DeptBioSci

W GREENLAND
Kobbefjord, BioBasis
Date: ___/___/2021
MART4C
GreenlandInstNatRes
AarhusUnl. DeptBioSci

W GREENLAND
Kobbefjord, BioBasis
Date: ___/___/2021
MART6C
GreenlandInstNatRes
AarhusUnl, DeptBioSci

W GREENLAND
Kobbefjord, BioBasis
Date: ___/___/2021
MARTSC
GreenlandInstNatRes
AarhusUni, DeptBioSci

Appendix 16 List with random numbers for microarthropod sampling

Year	Collection	Sample	X	Y
2019	1	A	1.5	5.5
2019	1	B	3	4
2019	1	C	5	2
2019	1	D	1	5
2019	2	A	2	1
2019	2	B	1	2.5
2019	2	C	3	7
2019	2	D	3.5	10
2019	3	A	4.5	7.5
2019	3	B	6	3.5
2019	3	C	4.5	6.5
2019	3	D	5	3.5
2020	1	A	1.5	4.5
2020	1	B	6.5	7.5
2020	1	C	5.5	7.5
2020	1	D	6.5	8.5
2020	2	A	10	6.5
2020	2	B	9.5	8
2020	2	C	6.5	2.5
2020	2	D	8.5	7
2020	3	A	9	5.5
2020	3	B	2.5	1
2020	3	C	1	7.5
2020	3	D	9	5
2021	1	A	2	9.5
2021	1	B	6	10
2021	1	C	0.5	4.5
2021	1	D	1.5	4
2021	2	A	1	5.5
2021	2	B	8.5	1
2021	2	C	0.5	6.5
2021	2	D	5	8.5
2021	3	A	2.5	1.5
2021	3	B	8.5	10
2021	3	C	1	4.5
2021	3	D	2.5	2
2022	1	A	0.5	3
2022	1	B	3	4.5
2022	1	C	2	4
2022	1	D	0.5	3.5
2022	2	A	6.5	1
2022	2	B	1	3.5
2022	2	C	6.5	0.5
2022	2	D	5.5	4
2022	3	A	4	9.5
2022	3	B	9.5	1
2022	3	C	3	8
2022	3	D	1.5	1